



06/22



ON THE  
IMMEDIATE TREATMENT  
OF  
STRICTURE OF THE URETHRA.





ON THE  
IMMEDIATE TREATMENT  
OF  
STRICTURE OF THE URETHRA,  
BY THE EMPLOYMENT OF THE  
"STRICTURE DILATOR."

BY  
BARNARD HOLT, F.R.C.S.,

SENIOR SURGEON TO THE WESTMINSTER HOSPITAL; LATE LECTURER ON SURGERY IN THE  
WESTMINSTER HOSPITAL SCHOOL OF MEDICINE, &c.

*Second Edition.*

LONDON:  
JOHN CHURCHILL & SONS, NEW BURLINGTON STREET.

---

MDCCLXIII.



## PREFACE TO THE FIRST EDITION.

THE greater portion of the following remarks has already appeared in the pages of the 'Medical Times and Gazette,' but in preparing them for re-publication I have appended some observations on the relative merits of the treatment I propose as contrasted with the various operations by incision; and have also added some cases to illustrate more fully some of the points on which I have insisted.

I have lately been able to examine the effects of forcible dilatation upon a stricture after the patient's death from another cause, —a point of much interest; and the description, with a drawing of the preparation, which was exhibited at the Pathological Society of

London, will be found at the end of the book.

The fatality that attends the uninterrupted course of a serious stricture, is but too certain. All practical surgeons are acquainted with the complications and suffering resulting from a disease, which, if taken in time, is *always amenable to treatment*. If, therefore, any means can be employed, by which all serious results can be obviated, and the treatment at the same time be made so simple as to be available by the majority of surgeons, it will, perhaps, be admitted that a step has been gained in the right direction. Hitherto I have not ventured to publish my experience of the plan I adopt, simply for the reason that it is injudicious to enunciate any new method of treatment which has not been subjected to numerous and repeated trials; but having now operated upon more than one hundred cases with unvaried success, not only in private but also in hospital practice, where every opportunity has been afforded

to the profession of witnessing the treatment and its results, I now feel justified in submitting that experience to the judgment of my professional brethren, and shall endeavour to frame such clear rules for their guidance, that, in any intelligent hands the operation may be as successful as it has been in my own. Up to the present time, this method of treatment, although adopted by some surgeons, has been mainly confined to myself, simply, I believe, from the fact that its general utility never having been published, it has not, as yet, been appreciated. Let me indulge a hope that, should I be fortunate enough to express myself so clearly as to be generally understood, the performance of the operation may become extended.

In conclusion, I would only deprecate the prejudices which exist in the minds of surgeons, against any novelty in the treatment of a common disorder, and request an unbiased perusal of the following pages.



## PREFACE TO THE SECOND EDITION.

THE progress of any new operation, more especially one for the cure of so serious a disease as stricture of the urethra, is deserving of record ; and I therefore gladly embrace the opportunity offered by the necessary preparation of a second edition of this book, of detailing my experience of the past eighteen months. During that period, I have treated some of the most formidable cases of stricture in all classes of society with unprecedented success ; and a perusal of these cases cannot but be interesting and instructive, and will further illustrate how little pain or inconvenience attends the new method of treatment, the results of which have been uniform,



and the danger *nil*. The apprehension that exists in the minds of some surgeons, that because the stricture is split, and no instrument is retained in the bladder, infiltration of urine must ensue, is perfectly fallacious. No single instance of such an event has occurred in any of the two hundred and fifty cases that I have now operated upon, and I believe it is all but impossible, the condensed tissue which surrounds an important stricture acting as a protective capsule, and preventing the escape of any urine. I have lately had two important modifications made in my instrument, which are described in the body of the work, and which will, I believe, remove every possibility of danger from its use.

The performance of the operation is becoming daily extended, both at home and abroad; and the cases lately published by Messrs. Fergusson, Paget, Smith, McNamara, Smily, &c., prove its success in other hands. The few additional cases are merely recorded as examples of many similar ones, and I trust

they will prove to the most sceptical, that difficulties which were hitherto almost insurmountable, are now in a very short time easily overcome; indeed, so strong is my faith in the operation, that I have no hesitation in affirming that *surgeons must adopt it*, since it is so simple, and has been hitherto proved so devoid of danger. I am quite aware that great prejudices have to be overcome; but I do trust, for the honour of our profession, that those prejudices will not be allowed to interfere with the progress of an operation capable of affording in so short a time such decided relief.

14 SAVILE ROW,

October, 1863.

out these kingdoms, and even awakened a surmise of the nature of these important truths in France and Germany. The pre-eminent abilities of Baillie, Abernethy, and Denman gradually moulded the practice of medicine, of surgery, and of midwifery in subjection to the Hunterian principles. A careful interpretation of Nature was the rigidly enforced rule. All rash, meddlesome, and coarse methods of treatment were denounced, the *vis medicatrix* was perpetually appealed to, *arte non vi* was constantly ejaculated. The late excellent surgeon, Mr. Lynn, an eminent pupil of Hunter, used, in his eccentric way, often to say, "*Nater*, gentlemen, *Nater* cures the disease."


Without doubt, this habitual subjection of the mind to the careful contemplation of natural processes was at the bottom of the marked improvement which, of late years, has taken place in every department of our art. It is a weakness of the human mind however, to oscillate from one extreme to the other. In medicine, from the extreme of heroic doses, and a licentious use of the lancet, the practical pendulum vibrated to homœopathic globules and a supinely expectant treatment. In midwifery, from excessive instrumentation, the accoucheur has occasionally fallen into a help-

less acquiescence in avoidable evils; and in surgery even an affected aversion to the knife has too often seduced the operator into a fatal procrastination.

The "native hue of resolution," which is nowhere more necessary than in the practice of surgery, has not, perhaps, in any class of affections been so injuriously "sicklied o'er," as in certain obstinate forms of stricture of the urethra. Mr. Abernethy's denunciations of all violence in catheterism as being inconsistent with the main purpose of the remedy—viz., the absorption of the material of the stricture—has had a greater effect than he probably contemplated. A timid and dilatory treatment has been the result, and in too many instances the supineness of the surgeon has permitted the case to drift to a fatal termination. It is a sad truth that "*Nater*" will not cure a chronic stricture.

My attention was early attracted to the prevailing defects in the treatment of strictures. I could not but observe the tediousness of the treatment by ordinary dilatation, occupying many months before an average instrument could be introduced into the bladder, and that even when the dilatation was accomplished, the contraction generally re-

FIG. 1.



turned, so that perpetual surgical care was required.

Being deeply impressed with the unsatisfactory nature of prevailing methods of curing these distressing maladies, about seven years ago I adopted a more energetic mode of treatment, and invited the notice of the profession to a new "Stricture Dilator." Its use was at first limited to simple dilatation, which was readily effected by graduated tubes passed between the blades without the withdrawal of the original instrument. Experience, however, soon showed me that, as a general rule, when dilatation was carried much beyond the degree produced by ordinary bougies, "stricture fever" was induced. I therefore determined, though with some apprehension as to the consequences, to *split the stricture* by passing the

largest-sized tube at once, and thus immediately to enlarge the contracted part of the canal, so that it might receive a catheter equal to the normal size of the urethra.

Fearing the effects of the urine being permitted to come in contact with the laceration thus occasioned, I kept a gum elastic catheter in the bladder, but as this measure gave rise to considerable irritation, I determined to content myself with simply splitting the stricture, drawing off the urine, and not again using the catheter till two days after the operation. After that interval, an instrument of the same diameter as that used at the time of the operation was again employed, and its use was continued—first, on alternate days, and, subsequently, at longer intervals. Experience has shown, indeed, that instances occur in which it is necessary to use a catheter one size smaller than that first passed after the operation, but such cases are exceptions.

The instrument by which this simple process is accomplished consists, as is shown in the drawing (Fig. 1) of two grooved blades fixed in a divided handle and containing between them a wire welded to their points, and on this wire a tube (Fig. 2) (which, when introduced between the blades corresponds to the natural calibre of

the urethra) is quickly passed, and thus ruptures or splits the obstruction. The simplicity of this apparatus is obvious to all, and my

FIG. 3.



experience of now above 250 cases proves that its use is unattended by any of those serious complications — viz., hæmorrhages, false passages, infiltration of urine, perinæal abscess, fistulæ, swelled testis, &c., &c. — which too often accompany the other operative processes devised for the relief of this malady. The forcible distention caused by the dilator affects the morbid obstruction only; the healthy portion of the canal not being disturbed, the slightly vascular character of the stricture deposition at the same time gives rise to but very inconsiderable bleeding, and perfectly obviates infiltration into the surrounding tissues.

Since the above was written, a very great improvement has been made

in the construction of the instrument, in my opinion so great as to get rid of every objection that could be previously urged against it. In the former instrument, fig 1, it was objected that there was no positive evidence that the instrument was in the bladder, and the objection was a valid one, although in the hands of an experienced surgeon, such a difficulty could hardly occur. This is now remedied by having the directing rod hollow, with an opening at the back of the curve of the dilator, so that when the stilette is removed (Fig. 3) the urine will escape in the same manner as it does when a catheter is used. But another objection was urged by some, viz., that it was possible for the tube to escape from between the blades of the dilator. This also as a *possibility* was true, but is now rendered quite impossible by the alteration that has been made in the handle; and as the dilator now acts as a catheter and the tube cannot be displaced, the only objection that can possibly be urged is, as to its introduction, and for this, each individual operator must be responsible.

The method of performing my operation may be described in a very few words. The permeability of the canal having been once satisfactorily ascertained, the size of the meatus of the urethra is to be gauged by passing into it



a sound that will conveniently fit, if it will admit a No. 10 use a No. 10 tube, if a No. 12 a No. 12 tube. For the No. 10 tube leave the screws in (Fig. 3) arranging them so that the tube will glide freely between the blades, but if the No. 12 tube is required remove the screws and the instrument is then set for that size. It is important to ascertain the proper measurement, so that the urethra may not be stretched beyond its natural calibre, for while the urethra of one person will admit No. 14, another will not admit more than No. 9.

The dilator having been previously well oiled, is to be introduced with the handle somewhat over the patient's left hip, and by keeping the convex portion gently pressing against the under part of the urethra, the point will glide along the upper portion until it is fairly beyond the triangular ligament, when, by bringing the handle to a right angle with the body, and gradually depressing it—but not so much as in the passage of an ordinary catheter—it will usually slip into the bladder; in fact, the same proceeding is to be adopted as in introducing a lithotrite for the purpose of crushing a calculus. The stilette in the perforated guiding rod is now to be withdrawn when the urine will flow, and being thus assured that the instrument is in the bladder, the Surgeon is next

to place the point of the tube he has previously selected, upon the wire between the blades, and thrust it as quickly as possible onwards to the end. By this means the stricture will be fairly split, and not dilated, the former effect being absolutely necessary to obtain the best results. The dilator should now be rotated to separate still further the sides of the rent, and then withdrawn; a catheter corresponding to the number of the tube being substituted, for the purpose of removing the urine. The catheter is then to be taken out, and the patient sent to bed, with directions to take, every four hours, for the first day and night, a mixture containing in each dose two grains of quinine and ten minims of the tincture of opium.

One caution is necessary to ensure the escape of the urine through the dilator, viz., that it should not be thrust so far into the bladder as to bring the opening in its curve against the posterior wall, since it would thereby be closed, and the passage of the urine would be prevented. Should this occur, let the dilator be partly withdrawn and the water will flow.

The facility with which this proceeding can be effected will of course depend upon the kind and number of the strictures, and the existence or otherwise of false passages, or

fistulæ in perinæo. The urine having been withdrawn, the patient does not require to pass water for some hours, and when compelled to do so, the stream is usually larger, and the urine passes with greater facility than before. On the second day from the operation, the same catheter should be gently introduced; but, if the patient complains of much scalding, it will be better to take one size less. This should be repeated every other day for a week, when the larger one may be substituted, and the patient be taught to pass his own instrument. Of course the time occupied in the after-treatment must vary with the nature of the case, and the more obstinate forms necessitate the employment of the catheter for some time, the intervals being gradually increased until it is not required to be used more than once in three, four, or six months, and in most instances, not more frequently than once a year. The bowels should be relieved by a dose of castor-oil taken early on the morning of the operation, and the patient should be directed not to pass water for two or three hours previously, in order—first, to facilitate the introduction of the dilator; and, secondly, to permit its free movement in the bladder.

A perusal of the following cases, which are extracted, as salient examples, from a long

series, and which were, for the most part, witnessed through their whole career by the students of the Westminster Hospital, and by those Surgeons who favour that institution by their attendance, will, I think, corroborate the points which I wish to establish, and, I hope, justify me in upholding as proved the following conclusions :—

1. That the operation is of the most simple kind, and that anyone who can pass a bougie through a difficult stricture is competent to perform it.

2. That it is not attended with hæmorrhage, infiltration of urine, abscess, or any serious local mischief.

3. That in the majority of instances the relief is immediate.

4. That the occurrence of rigors, or any other constitutional disturbance, is very rare, and the patient is seldom confined to bed longer than from twelve to twenty-four hours.

5. That the urethra is immediately made permeable by a catheter of full size, which may be ever afterwards passed at discretion.

6. That this method is available in every kind of stricture where a canula of any size can reach the bladder.

7. That when the after-treatment is judi-

cious and attentive, the full capacity of the passage is always maintained.

8. That in all cases of neglected after-treatment, the stricture yields again to this method more promptly than to any other.

9. That, it being impossible that any but the diseased tissue can be divided, the splitting of the stricture has a decided superiority over any cutting operation.

10. And, to sum up the great advantages in one proposition,—that the process is facile, speedy, prompt in its effects, and free from every danger, immediate or remote.

The course of general treatment will naturally vary, according to the kind of obstruction, the number of strictures, and the occasional complications of contracted bladder, enlarged prostate, fistulæ in perinæo, false passage, &c. In simple strictures, however narrow, the relief will be immediate, but in the more complex forms of these maladies, the size of the stream is not increased so directly as might have been anticipated from the immediate enlargement of the canal. Notwithstanding, however, that the size of the stream may for a short time remain somewhat restricted, the patient is able to empty his bladder much more quickly and effectually

than before, and has less frequent micturition. The limitation of the jet evidently depends upon the thickening of the surrounding textures, and swelling of the mucous lining; these morbid states speedily subside, and in a short space of time the patient can void his water in a normal manner.

### CASE I.

*Stricture of Twenty-five Years' Duration — Severe Constitutional Disturbance—Recurrence of "Stricture Fever" after each attempt to introduce a Catheter—Retention of Urine—Hæmorrhage—False Passage—Operation—Recovery.*

THOMAS W., aged 50, a labourer, of dissolute habits, was admitted into the Westminster Hospital under my care, on November 5, 1857. He had been the subject of stricture for twenty-five years, during which period the stream has been gradually contracting, and the urine is now passed *guttatim*. Five years since he was an in-patient at St. Thomas's Hospital, where, after considerable difficulty, a No. 1 catheter was introduced and retained, being replaced by others, in succession, until No. 6 could be passed, when he left. The contraction speedily recurred, and he was admitted into the Westminster Hospital as above.

He is now greatly emaciated, with brown and dry tongue; pulse 110; countenance pallid; appetite defective; abdomen tympanitic; nights restless; and there is a constant escape of fæces during the straining to evacuate the bladder, an effort which he is compelled to repeat every hour and a half. The house-surgeon having twice failed in passing a No. 1 catheter, each attempt being followed by syncope and subsequent rigors. I now saw the patient, and directed the nightly use of the warm bath, and prescribed salines with opium through the day, and castor-oil every morning. At the expiration of a week his general condition was sufficiently improved to justify the attempt to introduce a No. 1 catheter, when a stricture was detected five inches from the meatus, through which it was impossible to penetrate. Although the examination was conducted with the greatest gentleness, it was followed by a severe attack of "stricture fever," which was only relieved by the administration of opium. The urethra being irritable, and the patient much exhausted, another week was permitted to elapse, during which time quinine and stimulants were administered, but the second trial was likewise futile, and followed by precisely the same results as the first. After the lapse of a third week, and while

the patient was fully under the influence of opium, a third attempt was made, but with no better success. I therefore determined to place the patient thoroughly under the influence of chloroform, when after very considerable difficulty, a No. 1 catheter was introduced, and a pint of highly-offensive and turbid urine withdrawn. In the evening he had retention, which, after great difficulty, was relieved by a No. 1 gum catheter; there was considerable hæmorrhage, the patient losing from ten to twelve ounces of blood. He passed a restless night, and being unable to make water, the catheter was again had recourse to. (The house-surgeon having unfortunately omitted to retain the former one.) As it was found to be impracticable to re-introduce it, the man was again anæsthesiated, and another attempt made, but although the catheter passed to its whole length no urine followed, and it was evident that a false passage had been made. Opium and the warm bath were had recourse to, and in a few hours a small quantity of bloody urine escaped and continued to dribble away during the day and night.

When I saw the patient on the following morning, his tongue was dry and brown; pulse feeble and rapid; skin hot, dry, and



blanched ; and he complained of great tenderness upon pressure over the lower part of the abdomen ; he was immediately ordered calomel and opium every four hours, and a turpentine fomentation over the seat of pain. For the next few days he continued in a very precarious state, but the symptoms gradually yielded, and in ten days from the retention the urine was free from blood ; his health was, however, so much shattered by the accompaniments of the retention, that a month elapsed before any further instrumental attempt was made. Former experience having shown the impossibility of introducing any instrument excepting while under chloroform, he was once more anaesthetized, and the dilator with difficulty passed into the bladder ; the large-sized tube was then employed, the stricture fairly split, and the dilator removed. A No. 12 catheter was next inserted, and the urine withdrawn ; the bleeding was very trifling. Ordered a mixture of quinine and opium, two grains of the former and ten drops of the latter for a dose. In the evening he felt chilly and uncomfortable, but had no shivering ; the urine flowed with moderate ease, and was triflingly tinged with blood.

On the following day he was able to walk about the ward ; the urine came away in a

small but clear stream; the medicine was omitted, and his ordinary diet resumed.

On the second day after the operation, a No. 12 catheter entered the bladder without the least difficulty; the urethra was tender, but he did not experience so much pain as on former occasions; the after-treatment was continued at increasing intervals, and he shortly left the hospital capable of passing his own instrument.

The foregoing is a case of very considerable interest, as embracing many of the most important points in connection with stricture. The duration of the disease, the manner in which micturition was accomplished, the deplorable state of the patient's health, the supervention of "stricture fever" after every attempt at catheterism, the after establishment of false passage, and the profuse hæmorrhage, are all evidences of the serious nature of the case, which consequently presented a severe test to the adopted treatment. The history satisfactorily shows the immunity from fatal consequences, rapidity of recovery, and complete restoration of the urethra to its natural size, which are the great characteristics of the operation.

## CASE II.

*Stricture of Fourteen Years' Standing—Of a Dense Cartilaginous Character—Situated about the Triangular Ligament—Urine, which was Ammoniacal and Purulent, passed guttatim — Operation — Recovery.*

JOHN W., aged 45, a labourer, was admitted in December, 1857, suffering from stricture of the urethra, of fourteen years' duration. Seven years since, the urine having been for some weeks previously passed *guttatim*, he had retention, and applied at St. George's Hospital, where, after considerable difficulty, a catheter was introduced. He continued an in-patient until No. 4 could be passed, and then left, much improved both in health and manner of micturition. The stricture, however, gradually returned, and after several attacks of retention, and a generally 'increasing difficulty in relieving himself, he became an in-patient of the Westminster Hospital. At the time of his admission, the urine was passed every hour, with great straining, and frequently in drops; there was considerable hardness in the perinæum, and his health was much damaged, as evinced by emaciation, pallor, loss of appetite, sleepless nights, and general feverishness; the

urine was highly ammoniacal, and loaded with mucus and pus. A No. 1 silver catheter was attempted to be passed without success, there was complete obstruction at the triangular ligament, and although the point was firmly grasped it could not be made to penetrate through the stricture. The patient was kept in bed and the bowels regulated, and on the third day from his admission, another attempt was made in the most gentle manner, but with no better result. Considerable irritative fever followed the two trials, and it was only after the expiration of two months, during which time six endeavours were made, that a No. 1 catheter was passed. The stricture was of the cartilaginous variety, about an inch in length, and gave that peculiar grating to the catheter so specially characteristic of cartilaginous obstruction. A large quantity of highly-offensive urine was withdrawn, and the catheter was retained. On the following day, a larger size was substituted, and on March 4th, the patient having been placed under the influence of chloroform, the dilator was with considerable difficulty introduced, and the No. 12 tube immediately passed. A No. 12 catheter was then easily slipped in, and the urine withdrawn. On the following day there was no

febrile disturbance, and the patient declared he made water better than he had done for many years. On the succeeding day, or second after the operation, the urine was passed in a smaller stream, and with some scalding, consequently a No. 11 catheter was used, the urine being much less foetid than before. The stream continued small for a fortnight, but the urine was expelled in a much shorter time, and the frequency of micturition materially diminished; he now only required to relieve himself three times during the night, and the same during the day. No. 11 having been passed on alternate days, a No. 12 was introduced, and used every third and every fourth day in the manner described. Six weeks after the operation he left the hospital, making water in a perfectly natural manner, having been taught to pass a No. 12 with ease.

The foregoing, one of a class of strictures that offers the greatest impediment to the introduction of a catheter, is an excellent example of a formidable cartilaginous obstruction situated at a part of the urethra which, corresponding to its curve, gives greater trouble in the introduction of an instrument than any other. The patient had been the subject of a stricture for many years, his urine had for

months previous to his admission been passed *guttatim*, his clothes were constantly saturated, and his health was materially damaged by the hourly necessity for relief—an evil which depended upon the stricture, as proved by the large quantity of urine that was withdrawn when a catheter was introduced. All these difficulties were immediately overcome, and a large-sized catheter could be ever afterwards introduced with perfect facility. The offensive ammoniacal condition of the urine gradually abated, that fluid assumed a normal character, and the bladder being no longer irritated by its retention, the intervals between the times of micturition were prolonged. In a word, the patient was restored to complete comfort without having been detained in bed more than a day, and without experiencing a single unfavourable symptom ; indeed, he did not suffer in the same ratio as during the attempt to simply introduce a catheter. To those who are uninitiated in the treatment of stricture, it might appear that, by keeping a catheter in the bladder, and daily increasing the sizes, the same results would have been obtained, but experience proves this plan of treatment to be utterly futile, and that as soon as the catheter is removed, so soon does the stricture return.

## CASE III.

*Stricture at the Bulb of Twenty-eight Years' Standing—Frequent Attacks of Retention of Urine—Infiltration—Fistulæ in Perinæo and Scrotum—Enlargement of the Prostate Gland.*

H. R., a captain in the Navy, aged 58, consulted me after having for many years been suffering from the effects of stricture. He had had repeated attacks of retention of urine, and upon three occasions matter formed behind and in the scrotum, which eventually gave rise to fistulæ in those situations. At the time of his coming under my care, the greater portion of the urine escaped through these apertures, so that he was always compelled to make water in a sitting posture, or with the aid of an earthenware slipper held between the legs. His clothes were constantly saturated with these dribblings, and his health was materially damaged by constant straining, and the frequency with which he was compelled to make efforts to relieve the bladder. He had been under the care of many surgeons, both in London and elsewhere, but without obtaining amelioration of his sufferings. Having tried various sized and kinds of instruments, I was fortunate enough, at the expiration of three weeks, to get a No. 00 silver

catheter through the obstruction. This instrument was retained, and replaced by a larger size, until No. 3 could be passed, when the dilator was introduced, and the No. 12 tube immediately passed. The same after-treatment was adopted as in the preceding cases, and in three months from his first visit he was enabled to pass No. 12 without difficulty. The stream was good, the intervals much longer, and his health better than it had been for the previous seven years.

Another very similar case may be added—viz.: A gentleman, a solicitor by profession, was affected with a stricture of many years standing. He had for ten years experienced great difficulty in emptying his bladder, and the urine was loaded with mucus. Upon the first examination the catheter was arrested at a point three inches from the meatus; after some trouble it passed on to a second obstruction, near the membranous portion of the urethra, through which it was impossible at the time to penetrate. Upon subsequent trials, however, I got through the second obstruction, and came upon a third, nearer the bladder, which was eventually overcome. The dilator was at once introduced, and the three strictures split. A No. 12 catheter was imme-



diately passed in, and the urine withdrawn. Considerable pain was for a time experienced in this instance in expelling the urine, which was only effected with great straining. The cause however of this suffering was speedily apparent, for the patient passed three small calculi, which had formed behind the several obstructions, and were washed away when the urethra became enlarged.

#### CASE IV.

*Stricture in Front of the Triangular Ligament of Eighteen Years' Duration—Several Attacks of Retention—Infiltration and Abscess at the Root of the Penis—Operation—Subsequent Sloughing—Recovery after the Use of the Dilator.*

J. R., a man of sallow aspect, was admitted into the Westminster Hospital under my care for stricture. He has been the subject of stricture for eighteen years, during which period he has had several attacks of retention of urine. The urine for some months had been passed *guttatim*, and occasionally in the smallest stream; and a week prior to his admission, during a violent effort to relieve himself, he felt a sharp burning pain at the root of the penis, with slight relief to his

urgency, but no urine escaped *per urethram*. In the evening he had a distinct rigor, was feverish, and passed a restless night; and on the following morning noticed that the penis was swollen, of a dusky hue, and painful to the touch; his urine continued to escape in drops. The penis was fomented with warm water; but all his symptoms increasing in severity, at the expiration of a week he was admitted into the hospital. I detected infiltration of urine, and an abscess at the root of the penis, which was immediately opened; and, to prevent further infiltration and afford a free outlet for the urine, an incision was made into the urethra in the perinæum, posterior to the stricture, by thrusting in a sharp-pointed bistoury in front of the anus, with the back to the rectum, and cutting upwards and forwards during the straining of the patient. The urethra being opened, a gum elastic catheter was pushed through the wound into the bladder, and the urine thus permitted to escape. The catheter was firmly secured, and a poultice applied to the wound. In the course of a few days, the inflammation and swelling of the penis had subsided sufficiently to allow a No. 1 to be tried through the penis, but it was found impracticable to reach

the bladder. Upon a second attempt, the catheter went farther, but would not penetrate the entire canal; the third attempt was more successful, and, after considerable difficulty, the bladder was reached and the urine withdrawn. The catheter was allowed to remain until the following morning, when it was replaced by the dilator, and the stricture at once split by passing a No. 12 tube. My usual plan of treatment was adopted, and the urine withdrawn. On the following day, the patient expressed himself as having passed a tranquil night, without rigors, or even chilliness; the urine flowed in a moderate stream, and with but slight scalding.

On the third day from the operation, a No. 12 catheter was passed without difficulty, and this act was repeated on alternate days for a fortnight; the intervals were then gradually extended, and in six weeks from the operation, the perinæal opening having healed, he left the hospital, passing his own instrument, and making water in a full stream.

The above-recited case afforded another excellent opportunity of testing the efficiency of the treatment, and although undertaken at a time when the patient was hardly recovered from the effects of infiltration, it was attended

with signal success; he never had a bad symptom, and in six weeks left the hospital, so far cured that the stricture was entirely under his own control.

### CASE V.

*Stricture of Thirty Years' Standing—Fistulæ in Perinæo—Incontinence of Urine—Temporary Impermeability of Urethra—Subsequent Treatment by Small and Large Dilators—Recovery.*

J. R., aged 62, a well-formed and ordinarily a robust man, but now greatly emaciated, was admitted in 1857 under my care. He states he has been the subject of stricture thirty years, and attributes the origin of the disease to frequently-contracted gonorrhœas. Fifteen years since he was the subject of retention of urine, which was at that time relieved by the introduction of a very small catheter, but being unable to continue in the hospital where he obtained relief, the contraction speedily became worse, and an abscess subsequently formed in the perinæum, which, upon being opened, gave exit to pus and urine, the latter continuing to escape more freely through the perineal opening than through the urethra. He now became an inmate of a provincial hos-

pital, where every endeavour was made to get an instrument into the bladder without success, and he eventually left unrelieved, the urine continuing to dribble away, partly from the urethra and partly through the perineal opening, so that he was compelled to wear an apparatus specially made for the purpose of hindering the urine from saturating his clothes. No treatment having hitherto been of service, he came to London, and placed himself under the care of the late Mr. Charles Guthrie, who eventually requested me to admit him under my care. Upon examination, I found his statement to be true, and that the urine continually escaped. A No. 1 catheter was employed, but it was impossible to pass it beyond the membranous portion of the urethra, which appeared to be so encroached upon as to be almost obliterated. Various attempts were made from time to time without success, and he was, consequently, placed under the influence of chloroform, and a No. 1 catheter was forcibly passed into the bladder. This was permitted to remain, and on the following day was replaced by a gum elastic one. The patient did not suffer materially from the forced catheterism, but was relieved by the withdrawal of a large quantity of urine, and

in three days after passing the No. 1 silver catheter, I was enabled to introduce the dilator and split the stricture. The patient was then placed in bed, and the mixture of quinine and opium ordered. In the evening he was attacked with rigors, and experienced considerable scalding in passing his water, which he continued to do in a narrow stream, and in small quantities. On the day following he was better, but the urine did not run freely, and on the second day after the operation, the No. 12 catheter was, with some little difficulty, passed into the bladder. A small quantity of urine escaped, and upon this occasion the catheter was allowed to remain five minutes, and was then withdrawn. The same line of treatment was adopted for many alternate days, but with a No. 11 instead of a No. 12 catheter, the stricture being irritable, and not admitting the larger size. • A No. 12 was, however, eventually passed, and being taught to use the catheter for himself, the man returned to the country, making water in a full stream, and declaring himself perfectly cured. This patient was under treatment nine weeks.

Three cases of a precisely similar character have been under treatment, where inconti-

nence of urine, the result of over-distension of the bladder, was present, the patient suffering from the absorption of the deleterious properties of the retained urine, the effects of which were recognised by drowsiness, pallor, nervousness, and tingling or itching of the skin. In each case the patient has been subjected to the same treatment, and with similar success.

## CASE VI.

*Traumatic Stricture—Severe Hæmorrhage—Elastic and Recurrent after Dilatation—Operation—Recovery.*

J. H., aged 48, a master-builder, consulted me in June, 1860, in consequence of continued difficulty in passing his water. He stated that five years since, while walking along a roof, he suddenly fell across a beam. His perinæum was seriously bruised, and he experienced great pain in the urethra, from which a considerable quantity of blood escaped. He was taken home, and a Surgeon passed, with some difficulty, a gum elastic catheter, which was retained. The hæmorrhage gradually subsided, and the catheter was removed. He, however, soon became aware that his urine was passed with greater difficulty than formerly, and that he was compelled to strain

a good deal to get rid of the contents of the bladder. He again consulted a Surgeon, who passed a No. 2 catheter into the bladder, and gradually increased the sizes of his instrument until a No. 12 was attained to. The catheterism being discontinued, the stricture soon returned, and he again consulted the same gentleman, who recommenced the treatment, and soon arrived at No. 12. The convalescence was, however, of short duration, for, upon the treatment being discontinued, the contraction speedily recurred. Upon examination, I found no difficulty in passing No. 2, but the stricture would not admit a No. 4, and No. 3 was very tight. I therefore immediately pushed in the dilator, split the stricture, and afterwards passed a No. 12 with facility. This treatment was perfectly successful. There was never the least hesitation in introducing a No. 12 afterwards, and he continued to eject his water in a copious stream without straining and at the ordinary periods\*.

\* I may refer to two cases of severe traumatic stricture, treated by the same method by Mr. Heath, and published in the 'Lancet,' August 31, 1861, as confirming the applicability of this treatment to such forms of the disease.



replaced by No. 2. In the evening he had retention, but it was speedily relieved by the No. 1 catheter. He continued to attend every third day until No. 2 could be passed with some degree of facility, when the dilator was introduced and the stricture split, and No. 12 catheter immediately introduced. The urine then flowed through; and he walked home, having directions to remain quiet, and take the quinine and opium mixture.

On the second day from the operation a No. 12 was easily slipped into the bladder; he had had neither hæmorrhage nor shivering, and the utmost he complained of was a slight scalding. In ten days from the operation, this gentleman passed his own instrument, and in a fortnight returned to Paris, making water with ease, and using No. 12 without the least difficulty.

I saw this patient at the beginning of the present year. The No. 12 can now be passed with the greatest facility. He has never had either retention or difficulty since the operation, and may be considered well.

The cases which have been already narrated are selected as representing the most severe forms of stricture, and as being attended by complications which would materially militate

against any ordinary operation, and must have necessitated months of surgical attendance if the ordinary dilating plan had been selected. In every one of the recited cases the treatment was successful, and with but a little instruction the patients were enabled to maintain the after-control of the bladder, by occasionally passing a bougie. Although this treatment is specially advocated in the extreme forms of the disease, it is equally applicable in the milder or less complicated, more especially in cases where strictures are irritable and unyielding, and where catheterism is attended with difficulty and severe suffering.

In such instances the dilator gives immediate relief, and experience has proved that by splitting the stricture, tension is immediately overcome, and the passage of the catheter is made so facile, as to be deprived of any but the most trifling pain. The following cases will best illustrate this :—

### CASE IX.

J. R., aged 38, a barrister, consulted me in January, 1859, complaining of stricture of the urethra, from which he had suffered for the last eight years. He stated that he had had three attacks of gonorrhœa, the discharge ac-

companying each attack lingering for a considerable period, and necessitating the use of stimulating injections for its cure. During 1858 his symptoms became much aggravated: he was obliged to strain greatly to get rid of the contents of his bladder: the stream of urine which had become spiral, and occasionally forked, was very small, and his sufferings were further increased by the protrusion of piles during his efforts for relief. He had suffered from retention of urine, which upon one occasion followed a debauch, and on another resulted from inability to relieve himself during a railway journey.

Upon examination, I detected a stricture situated about the membranous portion of the urethra, through which a No. 2 catheter could, without material difficulty, be passed. The stricture was of short extent, apparently cartilaginous, and it grasped the catheter tightly. Having explained to the patient the *rationale* of the operation, he at once consented to its performance. The dilator was, therefore, immediately introduced, the stricture split, and a No. 12 catheter substituted for the purpose of withdrawing the contents of the bladder. There was the slightest possible bleeding, and he stated that the operation was not more severe than

the passing of a catheter. I directed him to remain at home during the afternoon, and to take the quinine and opium mixture.

This gentleman continued his attendance on alternate days for a fortnight; he never had the slightest difficulty in passing his water; the pain was very trifling, and a No. 12 catheter slipped into the bladder with the greatest ease. Being now taught to pass his own instrument, he was dismissed, cured.

#### CASE X.

J. J., aged 61, a gentleman, residing in Wales, consulted me in December, 1858, for stricture of many years' duration. He stated that during the previous year, the stream of urine had become materially decreased in size: that he was obliged to pass water every two hours, that the urine was highly ammoniacal and loaded with mucus: that the irritable state of his bladder compelled him to make water the instant he had the desire to do so: and that his clothes were constantly wetted, from his inability to eject the urine in a continuous stream. His rest was broken, and his health materially damaged by the absorption of the noxious properties of the retained urine. Upon examination, I detected a stricture at the triangular ligament,

which would only admit a No. 1 catheter. The obstruction was dense and unyielding, and about half an inch in length. As the patient was somewhat plethoric, I prescribed a purgative, and requested his attendance on the following day, at which visit the dilator was with slight difficulty introduced, and the stricture split. A No. 12 catheter was now substituted for the dilator, and the urine withdrawn. In this case, as in the former one, the cure was uninterrupted by a single unfavourable symptom; the patient was merely confined to the house one afternoon, and the same plan of after-treatment having been carried out, he, at the end of a fortnight, returned to Wales, capable of passing his own catheter.

I have had several letters from this gentleman since, in all of which he declares himself free from the slightest difficulty in passing his water or introducing his catheter.

### CASE XI.

*Obstinate Stricture of more than 20 years' duration, where great difficulty was experienced in passing any Instrument into the Bladder, and where the Operation was undertaken at the time the patient was suffering from Swollen Testis.*

ON Oct. 20, 1862, a gentleman was brought to my house by Mr. May, of Tottenham, suf-

fering from the effects of stricture of the urethra, for which disease he had 18 years previously consulted M. Ricord. The patient, an otherwise comparatively healthy man, æt. 43, informed me that for a considerable time he had experienced increasing difficulty in passing his urine, which came in a very small stream, with great straining, and that it was frequently only passed in drops: that he had had a series of rigors, and that his urine frequently contained a large quantity of tenacious mucus. Upon examining the urethra I detected a stricture at the bulb, through which I failed to pass a No. 1 catheter. There was slight bleeding, but apparently no further discomfort. On the 22nd, Mr. May wrote me that his patient was too ill to repeat his visit in consequence of severe rigors, succeeded by irritative fever. On the 27th he again consulted me, when I succeeded in passing the No. 1 through the obstruction at the bulb, but it was arrested at a second stricture about the membranous portion of the canal. The second attempt was followed by the same train of symptoms as the former, and I did not see him again until Nov. 6th, when I still found it quite impossible to pass the catheter through the second stricture. Other

trials were made during the remainder of the month, and it was only at the commencement of December, and after very considerable difficulty that I succeeded in passing a catheter into the bladder, this was, however, followed by swollen testis from the effects of which he was confined to the house for three weeks. On January 24th, although the testicle was still swollen, I succeeded in passing the catheter, which, under the circumstances, was retained for two hours, and upon its removal, I passed the dilator, splitting the strictures with the No. 12 tube, and substituting a No. 12 catheter, through which the contents of bladder escaped. The catheter was now removed and the usual medicine was prescribed. Although he had been previously the subject of rigors after manipulation, and frequently also without any attempt at catheterism, yet he had no shivering or unfavourable symptom of any kind, and his urine was passed in a large stream, but with some scalding. On the 26th the No. 12 catheter was introduced without the least difficulty, and the after-treatment was carried on for another fortnight, after which he succeeded in passing the catheter for himself, and he has continued to do so ever since, his health is now quite re-established,

and so far as the stricture is concerned he is his own surgeon.

The interest of this case consists in the difficulty that was experienced in getting a catheter into the bladder, the strictures being of such an unyielding character as to frustrate the many attempts that were made to overcome them. The constitutional symptoms were, prior to the operation, very marked, rigors following almost every attempt at catheterism, and hernia humoralis succeeding the passage of the instrument into the bladder; and yet, after the strictures were fairly split, no unfavourable symptom of any kind was experienced.

## CASE XII.

*Obstinate Stricture of many years' duration for which Mr. Syme's operation had been performed by himself—Internal Urethrotomy by another surgeon, and subsequently Mr. Syme's operation repeated—Return of the obstruction and inability of the patient to make water without the aid of a No. 2 Catheter—Immediate Dilatation—Cure.*

CAPTAIN T. was kindly sent to me by Mr. Cutler under the following circumstances:—Seventeen years since, he had stricture, which gradually became more and more contracted until, in 1853, he consulted Mr. Syme, who passed a No. 2 into the bladder, and sub-



sequently performed the cutting operation in the perinæum, which he advocates. The patient was confined to his bed for 16 days, and subsequently No. 9 could be introduced with moderate ease. The stricture, however, soon became again contracted, until only No. 3 would enter the bladder, when he again placed himself under the care of Mr. Syme, who dilated it to No. 9, the passage of bougies being attended with very great suffering. After leaving Mr. Syme, however, the stricture soon re-contracted, and internal urethrotomy was performed by a surgeon in England, after which operation he had infiltration of urine, and his life was placed in considerable jeopardy. The operation was not attended with the slightest benefit, and he was compelled to pass a No. 2 catheter every time he required to make water. His sufferings being now very great he had Mr. Syme's operation performed a second time, and a No. 9 catheter was again attained; this size, however, could not be maintained, and he soon went back to No. 8, No. 7, and, in four months, to No. 4, and then to No. 2, which, as before, he is now obliged to pass every time the bladder requires relief. It was under these circumstances that Mr. Cutler suggested his consulting me, and after ex-

plaining to him that I had never had so complicated a case, and that I could not answer for the result to be attained, I consented to perform my operation. On March 30, while under the influence of chloroform, I passed the dilator, and the No. 10 tube, but so much force was required to press the tube between the blades, that I was compelled to wrap a towel around it to protect my hand, and the tube was thrust down three or four times in order that the strictures might be effectually split; the urine was removed, and he took the usual mixture. In the evening I passed a No. 6 catheter without the slightest difficulty, and on the following morning he informed me that he had slept well, and had not experienced the least pain or inconvenience of any kind; this time I introduced a No. 7 as easily as I had done the No. 6, and removed the urine: he was desired to discontinue the medicine and take his ordinary diet. On the 22nd I found him perfectly well, and I introduced a No. 8 with ease, and in the afternoon he visited some friends at Northfleet, near Gravesend. On the 23rd No. 9 was passed, and on the 26th No. 10. On April the 1st he passed his own No. 9 without the least difficulty, and in a week afterwards he returned home, stating

that he could pass his water as well as he ever had done; of course directions were given him to continue the after-treatment.

The above case afforded an opportunity of testing the practicability and efficiency of the operation to the utmost. It belonged to a class for which the operation devised by Mr. Syme appeared specially applicable, and yet the disease returned within a short period of its performance, requiring two other operations, both of them perfectly useless, so far as protracted improvement was concerned, and of a character which not only placed the life of the patient in considerable danger, but necessitated his being confined to his bed: in the first instance for sixteen days, and in the second for six weeks. Whereas by the adoption of the operation I advocate he never went to bed at all, could pass water with perfect freedom from the time of the operation, and on the third morning returned to his friends at Northfleet, not having suffered the slightest inconvenience of any kind.

What explanation can be given of such an occurrence? It is so extraordinary that had not the case been seen by others a doubt might have arisen as to the accuracy of my statement. I believe the whole secret depends

upon the obstruction *only* being ruptured, and that no irritation or inflammation is excited by retaining a catheter in the bladder. It is impossible that any structure but the stricture can be ruptured; and when so ruptured it is left to heal of its own accord, the medium of union being dilated by the subsequent passage of bougies. But supposing that only a temporary advantage had been gained, and the operation had required to be repeated. The patient had not suffered anything, and had only been confined to the house a day and a half, and this more as a precaution than a necessity. It will be noticed that in this instance the catheter was passed in the evening, and again on the following day contrary to my usual custom; I shall again allude to this subject at the end of the book.

The following is an excellent example of the irritability caused by retaining a catheter in the bladder for the rapid cure of stricture; the case is well known, and was related by me in some observations at the Royal Medical and Chirurgical Society.

### CASE XIII.

MR. S., a Solicitor residing in the country consulted me on April 28, 1862, in conse-

quence of stricture of the urethra, from which he had been suffering for more than thirty years. He informed me that for more than twenty years he had experienced difficulty in passing his water, and that he had been under the care of two most experienced surgeons in London, one of whom had devoted himself specially to the consideration of such diseases, and that having with great difficulty passed a small gum elastic catheter into the bladder, the surgeon alluded to tied the instrument in, and on every alternate day replaced it with a larger one, until a size was attained that corresponded with the natural calibre of the urethra. He expressed himself as being gratified at the success which followed this plan of treatment, and fully expected the full size could be afterwards maintained; in a very short time, however, the stricture contracted so that he gradually receded to No. 2, which could only be introduced after considerable suffering, and with great perseverance. His health having been much damaged by the confinement necessary to carry out this treatment, and not having derived any permanent benefit from it, he declined, although he was urgently solicited to do so, to have the same method again employed. Under these circumstances he

was urged to permit internal urethotomy to be performed; but having a dread of the knife he likewise objected to that method, and after considerable suffering and frequent disappointment in the passage of even his small catheter, he determined upon the operation by splitting. So fearful was he of the consequences of the passage of a catheter being attempted, in consequence of frequent attacks of retention after the small instrument had been used, that he would not allow me to examine the urethra, but decided to come to town and have the operation completed without any previous examination. As he was a remarkably intelligent man, and could with great accuracy detail his symptoms, I consented to comply with his request, and on May 13, 1862, with very considerable difficulty, I passed a No. 2 silver catheter, and upon its removal introduced the dilator and split the strictures with the No. 10 tube. Although of a very nervous temperament he never gave the slightest expression of pain, either when the operation was performed, or when the No. 10 catheter was substituted for the dilator to remove the urine. Not being aware of his peculiar temperament, I prescribed the usual medicine of quinine and opium, as I was anxious, if possible, to pre-

vent his having any shivering, as under the previous treatment he had suffered seriously from rigors and their accompanying fever. The opium, however, acted as a poison and depressed him exceedingly, so that for some days his pupil was contracted, and he was generally drowsy and uncomfortable. In the evening he felt chilly, but it did not amount to shivering; he passed his water without the least difficulty, but with some scalding. On the 15th I passed the No. 10 with ease, and as the weather was fine and warm, I directed him to take a drive.

On the 17th the catheter was again passed, and from this time he has gone on uninterruptedly well, he visits me once in about five weeks, the No. 10 is maintained, he has never taken a dose of medicine and has gained more than a stone in weight, being now capable of going through a long day's shooting without the least fatigue.

The following case, although attended with the complication of infiltration of urine and perinæal and scrotal fistulæ, is a more striking example of the inutility of retaining a catheter in the bladder for the cure of stricture.

## CASE XIV.

GEORGE R., æt. 42, a coachman, was sent to me by Mr. Hemming, of Kimbolton, in May, 1863. He states that for twelve years he has been suffering from the effects of stricture, with increasing difficulty in micturition; that in 1854 he was an inmate of St. Thomas's Hospital. The surgeon under whose care he was placed having had considerable difficulty in passing a No. 1 catheter, this was retained and replaced from time to time by others, until a No. 11 could be easily passed, when, believing himself well, he returned to the country. But the stricture re-contracted at once, and in 1858 he became an in-patient of St. Bartholomew's; various attempts were made to pass a catheter, but without success, each attempt being followed by stricture fever. His health being much shattered, he returned to the country, passing his urine in the smallest stream, and occasionally by drops. In 1862, after great straining, the urethra gave way behind the stricture, and he became the subject of infiltration of urine; matter formed, and the abscess burst, leaving an opening in the



perinæum and scrotum, with considerable condensation in these situations, as well as above the root of the penis, corresponding to the symphysis pubis. Mr. Hemming with some difficulty passed a small catheter, and gradually dilated the strictures, (for there were two, one at the bulb, and a second at the membranous portion of the urethra), until a No. 8 could enter, but as soon as the catheters were discontinued, so did the contraction recur. Upon examination, I found a No. 4 catheter could be passed into the bladder, which in the hope of preventing urine burrowing in the scrotum, and so adding to his present mischief, I retained. The patient being desired to make his water entirely through the catheter, the different sinuses were laid into one, and he was desired to rub in mercurial ointment and iodine to the condensed mass on the scrotum and perinæum. The catheter in a few days becoming loose, was replaced by a larger one, and no urine was permitted to escape excepting through the catheter; the result of this was a considerable reduction in the swelling which gradually became soft, but the fistulæ did not heal. After two months continuance of this method the catheter was removed, and although an attempt was made on the follow-

ing day to pass the same catheter, it was found that the stricture had so contracted as only to admit one four sizes less; the urine likewise escaped through the unhealed opening in the perinæum. The catheter was again replaced, and the same treatment continued for another month, but with a like result—viz., that when the catheter was removed the stricture became as small as ever, and in a week would only admit a No. 4. Under these circumstances, I determined to try whether any benefit would arise from splitting the strictures, and having passed the dilator, split them with a No. 10 tube. No catheter was retained in the bladder. For the first four days there was slight weeping from the fistulæ, but at the expiration of a week they had entirely healed, and a No. 10 catheter could be passed with the greatest ease. The patient returned to the country, and the following is an extract of Mr. Hemming's letter to me:—"My dear Sir,—You have indeed achieved a victory in the case of George R., and I hope you will publish it in your next series of cases. It is doubly valuable from your long continued and pertinacious attempts to conquer the stricture and thickening by the old method before you tried your own; the failure which followed

your two months trial, and the immediate success which your own method brought about."

The above cases are excellent examples of the inutility of retaining a catheter in the bladder, although after its removal it may be reintroduced at short intervals, and the after-treatment be effectually carried out. I may say I have never seen a *bad* stricture benefitted by such treatment, and for the last twenty years I have abandoned such treatment as useless, the large size cannot be maintained, and the stricture very speedily re-contracts.

The following three cases are instances of obstinate unyielding strictures in which long continued catheterism was unsuccessful, but where the dilator relieved the patient in a very short period.

## CASE XV.

THE EARL OF ——— consulted me on Oct. 19, 1862, by the advice of Dr. ———, of Dublin. He stated that for many years he had suffered from stricture of the urethra, which had been occasionally benefitted by the passage of a bougie, but that for some time his infirmity had been getting worse, and that he could now only make water in the smallest stream,

with great straining, and that he was unable to retain his urine for more than an hour and a half. Upon examination, a stricture situated at the bulb was detected, through which a No. 1 catheter could be passed, but it was arrested at the membranous part, and I did not succeed in passing it into the bladder. There was slight bleeding, and he afterwards suffered severe scalding pain when the urine was evacuated.

The same treatment was persisted in until Oct. 30th, when a No. 2 was passed fairly into the bladder; this was repeated on Nov. 3rd, 5th, and 7th, and on the 10th, having at his own request been placed under chloroform, the strictures were split with the No. 12 tube, and the urine was removed with a No. 12 catheter. On the following morning he informed me that he had not suffered any inconvenience beyond being compelled to urinate every two hours, when the scalding was severe.

On the 12th, the scalding had somewhat subsided, and a No. 12 catheter was passed with great ease; this treatment was repeated on the 14th, 16th, 18th, 21st, 24th, and so gradually increasing the intervals, and on the 9th Dec. I saw him for the last time, as he could now pass his own No. 12 with perfect

ease. Lord ——— called<sup>1</sup> upon me in June of the present year, and informed me that he now made water as well as he ever remembers to have done, and that he could retain his urine for many hours; I passed the No. 12 catheter without the slightest impediment or pain.

### CASE XVI.

THE HON. COL. ———, who was invalided from China, in consequence of a severe stricture, consulted me in May, 1862. His history was very similar to the one just related, and he experienced about the same difficulty in voiding his urine. The bladder was, however, more irritable, and he was compelled to urinate the instant he had the desire to do so, so that while walking out he had to consider whether he should be in the neighbourhood of an urinal, at certain intervals, so that he might relieve himself without such special urgency. On May 10th, I tried to introduce a catheter, but failed, he became faint, and I desisted; this was repeated upon three occasions with a like result. On May 20th, however, I succeeded in passing a No. 1 catheter, and on May 31st, while under the influence of chloroform, the dilator was passed,

and the strictures were split with the No. 12 tube, the 12 catheter being afterwards substituted. In the evening he was comfortable, and did not complain of pain; there was no shivering or constitutional disturbance of any kind. On the following day he walked out, and on June 2nd, I again passed the No. 12 with great ease. The after-treatment was carried out at increasing intervals, and in July of the present year, after an interval of four months since a catheter had been passed, and fourteen months after the operation, I passed a No. 13 sound with great ease. This gentleman has never had the slightest difficulty in passing his water since the operation, was only confined to the house one afternoon, and is now in robust health.

#### CASE XVII.

SIR C. S—, an officer who had seen much service, consulted me at the request of Dr. Pettigrew. He had been for many years a martyr to stricture, and as he expressed it, had taken a house in Yorkshire to bury himself from society, his infirmity not permitting him to enjoy those pleasures which his position and age entitled him to expect. Very many trials had been

already made to effect an entrance into the bladder, but hitherto without success. He was compelled to make water every hour or hour and a half, and even during the intervals his water would escape involuntarily, so that his clothes were constantly saturated. On Dec. 12, 1861, I first tried to pass a No. 1, but like other Surgeons, was foiled in my attempts to pass the instrument into the bladder; there was a dense stricture at the bulb which defied the utmost patience. The 16th, 18th, 21st, 23rd, and 26th were expended in fruitless attempts. On the 28th, a No. 1 was passed into the bladder, and retained for half-an-hour. On the 30th, I again failed to effect an entrance. On January 1st, 3rd, and 5th, the catheter was passed, and each time into the bladder, and on January 8th, the operation was performed without chloroform, and the No. 12 catheter was immediately introduced; he hardly experienced the least pain and was out on the following day. The after-treatment was carried out, and in three weeks from the time of the operation, he felt himself so improved as to abandon his idea of going into Yorkshire, and as he was now offered another command provided he could leave England by the beginning of March I strongly advised him to accept it.

In the mean time he succeeded in passing his own instrument with perfect freedom, and by the time stated he left England to again assume a command abroad.

The last case was one of not only great importance, so far as the patient was concerned, but was an excellent example of the advantage derived from this particular treatment, as it would have been quite impossible for him to have done as he did had any other plan been adopted.

### CASE XVIII.

*Stricture of Long Standing Complicated with Disease of the Bladder, and probably of Ureters and Pelvis of the Kidney.*

J. S., æt. 35, a merchant, consulted me at the request of Dr. S., of Liverpool, in consequence of his suffering from long-continued incontinence of urine, which for a considerable period had necessitated his wearing an urinal.

When I first saw this gentleman in January of the present year, he informed me that for many years he had suffered from stricture and increasing difficulty in ejecting his water, and that two years since he suffered from total retention when he was in great agony, and no catheter could be passed. The urine, however,



eventually dribbled away, and it had continued to do so ever since, excoriating the penis and scrotum, and keeping him in continual discomfort. He also complained of great fulness about the lower part of the abdomen, and he walked with considerable difficulty.

The history of his case assured me he was suffering from retention of urine, although it had been escaping involuntarily during the last two years, and against his inclination, I persuaded him to permit me to pass a catheter. This, however, was not accomplished without great difficulty, a No.  $\frac{1}{2}$  being eventually introduced, when nearly two pints of highly offensive ammoniacal and purulent urine was removed. As it was evident that the bladder had lost its contractile power, the catheter was again had recourse to in the evening, and a further quantity was removed, and for the purpose of keeping the bladder as empty as possible, the catheter was passed three times on the following day. As the unyielding nature of his stricture rendered it difficult to, at all times, pass the catheter, I determined to operate at once, and consequently on January 20th (the No. 1 catheter having been previously retained for a short time), I passed the dilator and split the strictures with the No. 10 tube,

his meatus not admitting a larger-sized catheter than No. 10. In the evening he had a rigor and passed a restless night, and it was necessary to confine him to the house. On the following day, the catheter was had recourse to to remove the urine, and it could be passed with facility. On the second day from the operation, it was found that he could pass water in a small stream, although he was as yet not capable of emptying the bladder; the catheter was consequently more frequently passed than is usual, but as the great impediment had been overcome, he was able to pass a No. 6 gum-elastic for himself, so as to ensure the bladder being entirely emptied three times in the twenty-four hours. In a fortnight from the operation, he could very nearly eject the whole of his urine, and in a month he returned to Liverpool immensely improved in health, being capable of micturating in a full stream, and able to pass a No. 9 silver catheter without any difficulty. In June of the present year, this gentleman consulted me again, having suffered from an attack of inflammation of the bladder, but there was no difficulty in introducing the No. 9, although he had not done so for three weeks prior to my again seeing him.

## CASE XIX.

*Impassable Traumatic Stricture of Twelve Years' Standing—Retention requiring Operation—Cure of the Stricture by Holt's Dilator—With Clinical Remarks—Under the care of Mr. Holt and Mr. Heath. (From the Notes of Mr. Beadles, House Surgeon to the Westminster Hospital.)*

The following case has been already published in the 'Lancet' for June 13, 1863:—

"JOSEPH S., aged 34, was sent up to the Westminster Hospital on the 27th January, 1863, by Mr. Cubitt, of Stroud, to be under Mr. Holt's care for a severe stricture of the urethra.

*History.*—Twelve years ago the patient slipped off the roof of a house, and fell about nine feet astride the edge of a water butt. He lost a good deal of blood, but no urine passed, and repeated attempts at catheterism were made by three different practitioners, but without success. He then took some drops (tincture of muriate of iron?), and in twenty-four hours the urine began to dribble away, and has continued to do so ever since, the patient wearing a bladder for its reception. In December last he applied to Mr. Cubitt, who made

various unsuccessful attempts to introduce a catheter, and then sent him to Mr. Holt.

On admission the urine was constantly dribbling away, and his bladder was evidently much distended. The patient is continually straining to void urine, which causes him the greatest agony each time it passes. There is no scar in the perinæum, but the urethra in the region of the bulb can be felt to be thickened and very hard.

January 29th.—Mr. Holt made a careful attempt to introduce an instrument through the stricture, but was unable to do so.

February 1st.—A second attempt was made to pass a catheter, but unsuccessfully, and repeated on the following day with the same result.

3rd.—The patient became unable to pass any urine, and suffered much from the distension of the bladder. Mr. Beadles, the House Surgeon, therefore attempted to pass a catheter, but unsuccessfully, and then requested Mr. Heath, who was in the Hospital, to see the patient, and after very considerable difficulty he succeeded in getting a No. 1 silver catheter through the stricture and down to the prostate, where it grated against a calculus, and upon passing the finger into the rectum, a

number of calculi were felt imbedded in the prostate. It was found impossible to reach the bladder with the catheter, which was therefore withdrawn, after giving exit to a small quantity of glairy fluid (not urine). The patient subsequently passed a small quantity of urine with relief to the urgent symptoms. In the night these latter became aggravated, and he was in great pain from distension of his bladder. The house surgeon accordingly sent to Mr. Holt, who being unwell, and finding that Mr. Heath had seen the patient, requested him to visit the Hospital for him.

Mr. Heath put the patient under the influence of chloroform, and tried to introduce a catheter, and after several efforts, succeeded in introducing a No. 1 down to the prostate, where it again touched the calculus, but would not enter the bladder. Under these circumstances, Mr. Heath determined to make an incision into the perinæum, extract the prostatic calculi, and relieve the bladder. The patient being tied up for lithotomy, and the No. 1 catheter being held as a guide, Mr. Heath made an incision immediately above the anus, and cut down upon the catheter directly in front of the prostate. The finger introduced into the wound was passed carefully into the bladder,

and there discovered a large vesical calculus, in addition to numerous calculi lodged in the prostate gland. With lithotomy forceps, an attempt was made to extract the stone, but Mr. Heath found it was adherent to the left side of the bladder. A large quantity of urine escaped, and upon another attempt at removal the stone broke in the forceps, and was withdrawn piecemeal. A large number of small prostatic calculi were removed, and there was then found to be a quantity of calculous matter adherent to the coats of the bladder in various parts; this was scraped away with the scoop as far as possible, and removed. The patient was put to bed and had a dose of opium.

On examining the calculi carefully they were found to consist of two distinct formations, vesical and prostatic, the whole mass weighing 1084 grains, or 18 drams and 4 grains, and the prostatic calculi alone  $46\frac{1}{2}$  grains. The prostatic calculi averaged the size of barleycorns, and were of a dark brown colour. On fitting together the portions of the vesical stone there were found to be two distinct calculi. One composed of fusible calculus around a nucleus of uric acid, the pieces of which weighed 832 grains, and which had evidently been adherent to the coats of the bladder, and the other of

very remarkable shape, and weighing 104 grains, which consisted of a nucleus of uric acid very smooth, and of the size and shape of a sparrow's egg 399 grains, with a surrounding portion of phosphatic matter, more resembling an unci-form bone than anything else. These latter pieces came away in the forceps in the middle of the operation, but their source was very doubtful. The debris of phosphatic matter removed from the bladder weighed  $101\frac{1}{2}$  grains.

4th.—The patient is perfectly easy, the urine comes entirely by the wound in the perinæum.

10th.—There has been no pain at all in the bladder until to-day, when the patient complained of pain about the bladder, resembling that from which he suffered before the operation. Mr. Holt ordered him 40 drops of tincture of opium.

13th.—He has occasional fits of spasm, but is easy in the intervals.

19th.—The spasm and pain continuing, and the patient's health suffering from the constant distress, Mr. Holt had him put under chloroform, and proceeded to examine the bladder through the wound. After a careful examination, Mr. Holt succeeded in extracting two small pieces of stone apparently part of the outer coating of the former calculus. The care-

ful use of the finger showed the lining of the bladder to be in a very rough condition, but the great depth of the perinæum made it difficult to ascertain with certainty the existence of any more pieces of stone.

20th.—Is quite easy, and his pulse has come down from 120 to 84.

23rd.—General condition very satisfactory. The urine proving very alkaline, he was ordered dilute hydrochloric acid in infusion of buchu thrice a day.

26th.—Mr. Holt made an attempt to introduce a catheter per urethram, but did not succeed, and desisted on account of producing hæmorrhage. The urine all passes through the wound, but he can hold it for a short time, and then voids it into a bed pan.

March 2nd.—This morning part of the urine passed per urethram in a small stream, and was clear, and acid in reaction. Mr. Holt made another attempt to pass a catheter, and introduced it a little farther than on the last occasion, but did not reach the bladder ; patient very comfortable.

16th.—Mr. Holt succeeded to-day in introducing a No. 2 silver catheter into the bladder, and this was tied in. No urine passed by the perinæum during the day, and only a very



small quantity at night. He has no straining or pain in the bladder, and the urine is clear.

18th.—Mr. Holt removed the No. 2, and introduced a No. 3; this was withdrawn, but was again passed by the House Surgeon in the evening.

19th.—No 3 catheter was again tried, but without success, and a No. 2 was passed with a good deal of difficulty. The urine has for the last two days passed entirely through the wound in the perinæum, which is, however, of very small size.

21st.—The urine has passed in a very fine stream this morning. The stricture had evidently contracted, and no instrument could be passed.

24th.—Mr. Holt succeeded in passing No. 2.

28th.—The No. 2 was retained in the bladder until to-day, when Mr. Holt withdrew it, and succeeded in passing his dilator, when the stricture was immediately split up, and a No. 10 catheter passed. After the operation the patient had no bad symptoms, and only felt a little sore.

30th.—No. 9 passed without any difficulty.

April 1st.—No. 10 again passed, and this was done on alternate days until the 15th of April, when he was discharged cured.

In some clinical remarks upon this interesting case, Mr. Holt drew the attention of his pupils to its salient points:—First as regards the treatment of cases of injury to the perinæum, that an instrument should, if possible, be passed at once, and before any attempt at micturition is made, so as to avoid extravasation into the perinæum, and the probable formation of abscesses. Secondly, that if, as in this case, it was impossible to pass a catheter, the surgeon is bound to incise the perinæum upon a catheter or staff, introduced as far as possible, and having found the other end of the torn urethra, to pass the instrument into the bladder, and retain it until the integrity of the canal is restored. In this case it may be doubted whether the urethra was really torn across, although much injured, since when the man's bladder had most improperly been allowed to become overdistended, the urine dribbled away. With regard to the treatment pursued in the hospital, Mr. Holt said he was not at all surprised at having been foiled in his first attempt at catheterisation considering the severity of the case. And when retention occurred, the operation undertaken by Mr. Heath was no doubt the best with the knowledge of the existence

of prostatic calculi. The presence of a large vesical calculus was an unexpected complication which was successfully overcome; but the encrusting of the bladder with phosphatic matter was such a serious event that many surgeons declined to interfere with such deposits. Mr. Heath had, however, removed nearly all with the scoop, and although the mucous membrane must have been severely injured, the patient had scarcely any symptoms for a day or two. The subsequent return of symptoms, Mr. Holt attributed to the presence of the small portions of calculous matter which had eluded observation before, and their extraction gave immediate relief, bringing down the pulse in a remarkable manner. With regard to the after-treatment of the case, Mr. Holt observed that, in consequence of the extreme difficulty which was experienced in getting any instrument into the bladder, he departed from his usual practice and tied the catheter in, but after the withdrawal, even for a day, the contraction immediately recurred, although the catheter was passed more frequently than usual. So he was eventually obliged to pass the dilator immediately the catheter was removed, and split the stricture at once."

This case served to illustrate admirably the result of the operation, as the man has gone down into the country able to take a No. 10 easily, and if the instrument were passed every few months would probably have no further trouble with his urinary organs.

## CASE XX.

*Obstinate Stricture—Abscess in the Perinæum—Fistula in Perinæo, &c.*

ABBOTT L., a pallid, unhealthy looking man, æt. 40, was admitted under Mr. Holt in June 1863. He stated that for many years he had experienced difficulty in passing his urine, and that four years since he had an attack of retention, which was, however, relieved by the passage of a catheter. Eighteen months since the stream became very much diminished, and he shortly afterwards placed himself under Mr. Wood's care in King's College Hospital. Mr. Wood succeeded in passing a No. 3 catheter upon two occasions, the difficulty being upon each sufficiently great to necessitate the administration of chloroform; in consequence, however, of some business that required his attention he was compelled to leave the hospital without deriving the full benefit of Mr.

Wood's treatment, and until the following April he neglected applying for any advice. At that time, he experienced greater difficulty than ever, and noticed a swelling in the perinæum, and that his urine contained pus.

On admission he informed the house-surgeon that he had not passed water for sixteen hours. He was in great agony, and the house-surgeon endeavoured to pass a catheter for his relief, but not succeeding, he placed him in a warm bath, and prescribed opium, under the influence of which he passed a small quantity of urine. On the following morning, the patient was seen by Mr. Holt, who immediately made an opening into the swelling in the perinæum, which gave exit to about an ounce of pus, and as the incision was made to extend into the urethra, the urine was afterwards freely evacuated through the wound. The patient was in a very low state—partly from his sufferings connected with the retention of urine and partly from bronchitis, from which he had been for some days under treatment. On the following morning he was much relieved; he had passed a quiet night, slept well, and his urine was freely evacuated through the wound in the perinæum. Mr. Holt considering the then unsatisfactory state of the

patient, and that the bladder could be emptied, did not deem it advisable to attempt the passage of a catheter, hoping that as the urine no longer came in contact with the stricture, it would gradually become less irritable, and more amenable to treatment. The patient continued to pass his urine entirely through the wound for the next three weeks, when, as the opening was becoming small, Mr. Holt tried to pass a catheter, but without success; the parts bled freely, but he was not otherwise inconvenienced. Attempts were now continued at various intervals without, however, the catheter reaching the bladder, and upon two occasions retention again ensued, so that the wound in the perinæum was obliged to be re-opened, for although the catheter had latterly passed through the front stricture, it would not reach the bladder. Under these circumstances a director was passed through the perinæal opening into the bladder, and upon this director the point of the catheter was pressed forwards, so that by this means the bladder was eventually reached. The catheter was now tied in, in the hope that the urethra would speedily become sufficiently enlarged to permit the ready entrance of the dilator, but as this did not occur, the dilator, in a few days after-

wards, was passed upon the director in the same way that the catheter had been, and the strictures were immediately split with the No. 10 tube. From that time the patient had no difficulty in voiding his urine, and he shortly left the hospital with his urethra admitting a No. 10 easily.

The surgeon is occasionally consulted in cases where the patient frequently suffers from retention of urine, although a moderately large-sized catheter can be passed. The following two cases are good examples of the benefit derived from at once enlarging the urethra and destroying the source of irritation.

### CASE XXI.

CAPTAIN P——, a patient of Mr. Miller's, of King Street, St. James', consulted me by that gentleman's advice, in consequence of his continually suffering from retention of urine, which compelled him always daily, and sometimes more frequently, to pass his own No. 7 catheter.

He informed me that some years since he had suffered from stricture, and for which he had been successfully treated, but that latterly, although he could pass his urine in a stream,

and easily introduce a No. 7 catheter, yet that daily he had retention of urine, for the removal of which he was compelled to resort to the catheter. He was a very abstemious liver, but had for a long time suffered from chronic rheumatism. Upon examining his urethra, I found no difficulty in introducing a No. 7, but it would not admit a No. 9, the obstruction being situated about the membranous portion of the canal. Believing this to be a case of irritable stricture, not easily dilated, and where spasm was sufficiently strong to cause retention where No. 7 only could be passed, but where probably such would not be the result if No. 12 could enter, I persuaded him to have the operation performed, and on Jan. 29, 1862, without the aid of chloroform, I introduced the dilator, and split the obstruction with the No. 12 tube. A No. 12 catheter replaced the dilator to empty the bladder, after which it was withdrawn. On the following day he informed me he had not suffered from the operation, and that his water had been passed less frequently, and in a full stream. On the 31st, the No. 12 was again introduced without any difficulty, and so the after-treatment was continued on alternate days without any retention, until Feb. 12th,



when the urine, being loaded with lithic acid, he found himself unable to relieve the bladder without the aid of the catheter, and a No. 11 was easily passed. From this time he continued uninterruptedly well, daily gaining strength, and capable of passing his own instrument without the least difficulty. This gentleman called upon me in July of the present year, and I passed the largest instrument without the slightest hesitation. He informs me he has never suffered from retention since he left London, and that he considers himself in perfect health.

## CASE XXII.

MR. H——, a patient of Mr. Alfred Heales, of Luton, consulted me under similar circumstances, and although in his case a fair sized instrument could be passed, yet after any excess or undue exertion, he was troubled with retention of urine, and was obliged to have recourse to the catheter. This patient remained under my care for a very short time; there was but little difficulty in passing a No. 5, so that I introduced the dilator at once and enlarged the urethra, so that it would at once admit a catheter of the natural size of

the canal. There was not the slightest complication or suffering, and he returned to the country in a fortnight, not having had retention since the operation, and with the urethra so enlarged that it would admit a large-sized catheter. I have since received a letter from Mr. Heale, who says "I have just seen H——, whose stricture you were good enough to rupture ; it is, I think he said, three weeks since his return, he expressed himself as much relieved, and thankful that he has not since had an arrest of urine. To-day I passed No. 10 easily ; I have no doubt of the efficacy of the plan of treatment, and I shall have no hesitation in sending you a patient another time."

Were I so disposed I could continue relating cases of extreme interest, and of great gravity, but as they would be a mere multiplication, acknowledging a simple principle, I shall content myself with recording the more interesting example of a most resisting and unyielding stricture in a patient whom Mr. Arnott was kind enough to place under my care.

## CASE XXIII.

COL. C——, a very stout, heavy man, about 50 years of age, had suffered from stricture for many years; he had consulted all the celebrated surgeons of the day, and his urethra would now only admit a No. 6 catheter, which was passed with considerable difficulty, and was so tightly grasped as to require considerable force to remove it. The stricture after a time would yield so as to admit a No. 7, and even a No. 8 had been pressed in, but no benefit was derived from this treatment, and he continued to pass water in as small a stream and as frequently after the catheter had been passed as before; in fact, the dilatation plan could not be carried out. I might mention that internal urethrotomy had been previously performed, and that the urethra had been cauterised after the most approved fashion. He had several times suffered from retention of urine, and severe rigors followed almost every attempt to enlarge the diameter of the urethra. The prospect was not a very pleasing one, as all other plans had failed in affording the required relief, and as severe constitutional symptoms succeeded each passage of a larger instrument

than usual, I could not ensure the patient against the probability of their following my operation, although I had, in many instances, found no inconvenience succeeding to the rupture of the stricture, although severe rigors had supervened upon catheterism. On June 28th, 1862, I performed my operation in the presence of Mr. Arnott, who was kind enough to accompany me. Although the dilator could be easily passed into the bladder, I never was compelled to use so much force for the rupture of the stricture, excepting in the case of Capt. T——, who had been operated upon by Mr. Syme. The Colonel bore the operation without a murmur, and a No. 10 catheter was afterwards passed, but very little blood escaped. In the evening he had a violent rigor, accompanied by vomiting, this succeeding to his passing water, which scalded him severely; his skin was hot; pulse 120; and he complained of pain in his head and back. Some saline medicine, with opium, was administered, and in the morning he was more comfortable, as the skin had freely perspired; the stream of urine was not improved, but he had passed it in small quantities. A No. 9 catheter was easily passed, and about 8 oz. of urine were removed; he remained for the

next few days in a very languid state, and as the rigor had been severe, the catheter was not passed until July 3rd, when a No. 10 was attempted; it was firmly grasped, and did not quite enter the bladder. In the evening he had another rigor, for which the same medicine was prescribed; the urine continued to be passed in a small stream, and he was very weak. On the following day he was somewhat better, tonic medicines were now prescribed, and wine fairly administered, and under this treatment he continued steadily to advance, although he suffered from a kind of bastard ague, even when the catheter had not been passed. On July 23rd, another attempt was made with No. 9, which passed readily into the bladder, and was not succeeded by rigor, and on the 25th No. 10 was again passed but it did not quite enter the bladder, and the attempt was again succeeded by rigor. No further attempt was now made until Aug. 2nd, when a No. 11 passed easily and smoothly into the bladder, and was not in the least degree grasped, when it became necessary to remove it. On the 4th, the same instrument was had recourse to; it entered the bladder, and was not followed by any unfavourable symptom. He was now taught to

pass his own bougie, which he did without difficulty, and in a few days afterwards went to the sea side to recruit his health. This gentleman has remained perfectly well ever since; he has greatly improved in health, and passes his No. 11 without any difficulty.

To prove that the treatment I advocate is equally applicable in cases of stricture of the female urethra, I will record a case which has been already published by Mr. Curling in the 'Lancet' of June 14th, 1862.

#### CASE XXIV.

*A Case of Close Stricture of the Urethra in the Female,  
Cured by Forcible Dilatation.*

ANN W——, aged 39, married, was admitted on the 17th of March, 1862. She is a pale, delicate-looking woman, a native of Woolwich, and states that she has been married twenty years and has had two children. She was married eight years before the birth of her first child; the labour was natural, and the child lived. Four years afterwards she gave birth to a second child, but she was this time attended by a midwife, who, she states, was intoxicated at the time, and "forced the labour." She is under the impression that the midwife ruptured

the membranes, and used great violence in extracting the child, causing her to "scream with the agony." There was no one present at the time. The child was born alive, but died soon after its birth, its insteps, hands, and throat having been severely lacerated by the midwife's rough usage. She had immediately afterwards great cutting pain on passing urine, which, she says, contained a large quantity of blood. At the end of three days an abscess formed in the neighbourhood of the urethra, producing almost complete retention. She was confined to her bed for four months, her medical attendant ordering her lotions and medicine, but she refused to have any instrument passed. Soon afterwards she left England, and sailed to the West Indies in company with her husband, where she remained eight years. She continued to have more and more difficulty in passing her urine, which came away in a very fine stream and in small quantities. Her bladder became irritable, and she sometimes had complete retention, which she was in the habit of relieving by hot fomentations, &c. In March, 1862, she was sounded by a surgeon in the country, and sent to Mr. Curling, under the impression that she was suffering from stone in the bladder.

On the 17th of March she was admitted by

Mr. Curling, suffering from great difficulty in passing urine, attended with a cutting pain. The urine contained a considerable quantity of mucus, but was acid.

In the ward an examination was made, when the meatus urinarius was found displaced and hidden by several fleshy-looking, projecting bodies; and, owing to the unfavourable light and bad position, no instrument was introduced. A warm bath was ordered, and half a grain of morphia at bed-time, to be followed by castor oil on the next morning.

On the 22nd, she was brought into the operating theatre, and examined on the table, when the meatus was found behind an excrescence on the left side. It was extremely small, and the urethra was so contracted that a small probe was with difficulty introduced. The examination was followed by great pain. An enema of gruel, with forty minims of tincture of opium, was administered, and she passed a good night.

On the 2nd of April, as she was rather low, four ounces of wine were added to her ordinary diet.

On the 4th, she was again brought into the theatre; chloroform being administered, Holt's dilator was introduced, and the stricture split



by the passage of a middle-sized tube. This was withdrawn, and a short No. 6 elastic gum catheter was passed into the bladder. This was tied in, and remained for four days, an anodyne and a warm bath at night being administered, which gave great relief to the dragging pains of which she then complained.

On the 7th, Mr. Curling introduced No. 9 elastic gum catheter, and next day No. 11 was got in by the house surgeon ; but the day after, as it was causing some distress, it was discontinued for two days.

11th.—She was not so well ; had a little fever, with loss of appetite, headache, and sickness. She was ordered a saline effervescing mixture, which stopped the sickness.

Three days after, a No. 9 catheter was ordered to be passed daily. To continue the effervescing mixture, as she was still a little weak and sick, with milk, beef-tea, and light pudding.

The following day she passed a No. 7 catheter for herself, and was better in health.

17th.—Urine a little loaded with mucus, and very acid. Ordered by Mr. Curling twenty grains of bicarbonate of potash, with thirty minims of tincture of henbane, thrice a day.

Next day she said she was much better and the bladder was less irritable.

Being anxious to join her husband, she was discharged on the 19th. She left the hospital considerably relieved, was able to pass her urine in a full stream, and had lost a great deal of her former pain. She promised to continue passing a No. 8 catheter, and expressed herself as much relieved by, and grateful for, the treatment she had received during her stay in the hospital.

The following are some clinical remarks on the case made by Mr. Curling:—

“The urethra of the female, being short, simple, and seldom affected with inflammation, rarely becomes the seat of stricture. A few cases have fallen under my notice. You may know that the worst form of stricture in the male arises from injuries, lacerations, or contusions. In all cases of stricture in the female which I have seen, the disease has arisen from injury of the parts in labour. In this case we have a clear history of a badly-managed labour by a drunken midwife, who used great violence in extracting the child. On voiding urine afterwards the patient passed blood and suffered great pain, and an abscess formed in the neighbourhood, which produced almost complete re-

tention. Difficulty in micturition was experienced for eight years; and when she was admitted into the hospital the urine was passed in a very fine stream, and she was suffering from chronic cystitis. The state of the parts and the displacement of the meatus gave every evidence of laceration and injury at some former period, and only a probe could be passed into the bladder. Some years ago I met with even a worse case of stricture than this, arising from injury in labour twenty-eight years before. The stricture was so small that I failed in getting an instrument into the bladder; and as the woman was suffering from retention, I had to puncture the bladder at the seat of stricture, and drew off from thirty to forty ounces of urine. The stricture was treated afterwards by dilatation. In the case of Ann W——, two methods of treatment presented themselves: by incision and by forcible dilatation. It was impossible to pass any instrument of the size required for making internal incision; and if this were practicable the operation would be attended with the risk of destroying the power of retaining urine. We see this after incising the urethra in the female for the extraction of stone. I determined, therefore, on trying forcible dilatation by Holt's

instrument. My experience of this treatment is very slight; but in cases where it has been employed, the urethra, contrary to my expectations, has borne forcible stretching without injurious results, such as hæmorrhage, infiltration of urine, abscess, or any serious local mischief. In this case, after forcible dilatation, I was enabled in a short time to pass a good-sized instrument into the bladder, and, considering the severity of the stricture, a cure was rapidly accomplished; and as the patient has learnt to pass instruments for herself, we need not be anxious about a return of the disease. There is every reason to expect that she will be permanently cured."

The foregoing cases have been briefly described, and no allusion or comparison has been made to the various kinds of treatment already published. Their perusal must have satisfied the most sceptical that, in every instance where any kind of instrument can be passed into the bladder, the urethra may, by the mechanical effect of the dilator, be immediately enlarged to its natural size; and that while in slight cases this enlargement is effected without suffering, so as to render the inhalation of chloroform unnecessary, yet in

the more severe forms that agent facilitates the operation, and secures a perfect immunity from pain. It is quite true that if the after-treatment is not attended to, the stricture will sooner or later recur, but as the patient by passing his own instrument has the control of the bladder, it can only return as a consequence of culpable neglect. The immunity from accidents having been already proved, the surgeon need have no fear of those serious results which so frequently accompany any cutting operation, and it is no less extraordinary than true that whereas rigors ordinarily supervene after rapid dilatation, they form the exception where the stricture has been fairly split.

It may be asked, what advantages accrue from the performance of this operation, when it is necessary that the after-treatment should be continued for a variable period of time: and why the ordinary plan of dilatation should be abandoned? My reply is—1st, that by its performance a large-sized catheter can be immediately passed, and the patient be speedily taught to use his own instrument; 2nd, that the patient avoids all the suffering incidental to gradual dilatation, and the frequent disappointment of not being enabled to increase the

sizes of the instrument ; 3dly, that it secures all the advantages that can accrue from the performance of any operation, and with much less danger to the patient than by any cutting operation ; and, lastly, that the period of recovery is shorter, it does not necessitate confinement to bed, and the patient is saved the expense of being constantly under the hands of his surgeon.

These observations apply to those cases where dilatation can be continued in the ordinary manner ; but all practical surgeons know that there are a large number of cases where dilatation is ineffectual in advancing beyond a certain gauge ; and here some operation must be performed, if the patient is not to continue a surgical annuity.

I am fully aware that in advocating a plan of treatment by which the stricture is split or torn, prejudices have to be overcome, from the fears which naturally arise as to the extent of the rupture, and the consequences, not only to the urethra, but also to the immediately investing structures. What may be the precise limit of this rupture in the living body I have not as yet had an opportunity of ascertaining, the treatment having been attended hitherto with unvarying success ; and when already

more than 250 cases have been operated upon by myself, in addition to those where the operation has been undertaken by other surgeons, in all of which the treatment has been most efficacious, it is not too much to hope that the danger must be very limited, when compared with any of the cutting operations.

What are the advantages of incising the urethra in those obstinate forms of stricture which necessitate some operation? Is it a more simple operation, less dangerous, and possessing any peculiar advantages in reference to a recurrence of the disease? I unhesitatingly state that there are no such advantages, but that, on the contrary, the operation is more difficult, and requires considerable experience for its performance. The operator in only a very limited number of cases, can insure the limitation of his incision to the strictured part, and for its performance it is necessary that some instrument should reach the bladder. Do the records of these operations give that degree of success which has been obtained by the operation I advocate: and setting aside the complications of hæmorrhage, infiltration of urine, abscess, &c., which so very frequently ensue, do they offer any better chance of permanent cure than the divi-

sion of stricture by rupture? Is not their after-treatment more tedious, the patient being confined to bed with a retained catheter until the greater portion of the wound has healed; and when this has united, do they not necessitate the rigid enforcement of the after-passage of the catheter, to ensure a prevention of the contraction?

The only difference between an incised wound and a rent, is that the former, if the edges are left approximated, heals much more quickly than the latter, and so more speedily induces a recurrence of the contraction. It is by stretching the uniting medium of either incision or rent that the cure is to be accomplished, just in the same manner as the orthopædic surgeon extends the soft union of a divided tendon.

Mr. Syme, with that ingenuity and surgical skill which have so long placed him in the highest rank of our profession, devoted his attention to the treatment of this most important disease, and devised a plan which, at the time of its publication, offered, in the more obstinate forms of stricture, the best chances of success, more especially when the contraction was situated in front of the anus. But even in the hands of this scientific surgeon, the



operation has not been without its attendant evil consequences, and in those of surgeons of less experience, has frequently terminated fatally,

Knowing the dangers incidental to perineal section, various surgeons, both at home and abroad, have endeavoured to substitute an internal for an external incision, and various instruments have been invented for the purpose of incising the stricture from within ; but with all there is the serious difficulty of not knowing with accuracy the exact structure which is being divided, and as the cutting portion of the instrument is usually *set* to divide a stricture, of the extent of which the surgeon is ignorant, it has been frequently found to cut either beyond the thickness of the stricture, or to have divided a portion of the urethra, either in front of or behind the obstruction ; hence, the frequent accompaniments of serious hæmorrhage, infiltration of urine, and abscess.

To accomplish either operation, it is necessary that some instrument should pass through the stricture ; and when that can be effected, I maintain that no cutting operation is either necessary or justifiable, and simply from the fact, that where any instrument can be passed, there the dilator can enter, and by the intro-

duction of the tube (which, with the dilator, should only represent the natural diameter of the urethra), we can split or tear that which is the seat of obstruction; and as far as I have yet had an opportunity of judging, the obstruction only. There is also another material advantage which attaches to splitting in preference to cutting—viz., that the surgeon is not obliged to keep any instrument in the bladder, but the patient passes his urine in a natural manner ever afterwards, the catheter being introduced at first on alternate days, and afterwards at longer intervals, until the canal is so far widened that the occasional passage of the bougie will maintain its calibre. While, however, I am unquestionably of opinion, from the large number of cases that I have operated upon, that the operation by rupture is the most feasible, the easiest to perform, and the least dangerous, I do not place it before the profession as a means of cure, unless the after-passage of the bougie is faithfully carried out. We know not at present of any means by which a stricture can be cured without such after-treatment, any more than the orthopædic surgeon knows of any means of curing deformities without the application of some extending apparatus after a tendon has been

divided. Our extending, or rather distending, means consists in the introduction of the catheter, or sound, and I care not whether the operation be by incision or splitting, the disease will recur if the contracted part of the canal is not kept dilated for a given time, such period varying with the nature and number of the strictures, the irritability of the patient, and the complications of false passage, fistulæ in perinæo, &c., &c., which may exist.

The advocates for internal incision affirm, that inasmuch as the knife is applied solely to that portion of the urethra which constitutes the obstruction, they divide the stricture only ; I must, however, express my doubt as to the accuracy of this statement, and record my belief that it is quite impossible to determine, at six or seven inches from the meatus, which side of the urethra is thickened ; the sound, or cutting instrument, is firmly grasped, and unless the seat of thickening, or deposit, is always the same, it will be quite impossible, from the sensation conveyed to the surgeon, to know which part to incise.

The numerous preparations in the different museums prove that, although the under part of the urethra may be the most frequently diseased, yet that the exceptions are so numer-

ous as practically to render that fact of no value, and that where the obstruction is of the obstinate character met with in the cases I have detailed, the thickening may be taken to be tolerably circumferential.

Suppose it were otherwise, and that the upper segment of the urethra were not involved, what should prevent the onward passage of a catheter when it has entered the contraction, since the upper part, if normal, would certainly yield, and permit of its passing to the bladder? The truth is, we have no security that, in the employment of a cutting instrument, our incisions are confined to the stricture alone, or even that the weakest part of the urethra may not be divided, and thus give rise to infiltration of urine and abscess, an occurrence by no means unfrequent where such means are employed. Hitherto, any treatment beyond ordinary dilatation has been considered applicable only to cases where the most severe complications are met with, and where the difficulties have been such as to compel the patient to place himself under the care of a surgeon who has great experience in the treatment of such cases; but the ordinary circumstances of cases, as they are usually presented to us, offer no obstacle to their immediate sub-

jection to the practice I advocate. As a general rule, a patient seeks professional aid when a No. 3 or No. 4 can be introduced; and here, no matter what may be the exact pathological change, the stricture may be immediately split, the catheter being afterwards passed upon three or four occasions by the surgeon, when the further treatment of the case may usually be transferred to the patient, who is now capable of passing his own instrument. The three following cases will best illustrate this:—

J. W., aged 45., consulted me in January, 1858, in consequence of difficulty in micturition, the result of stricture at about six inches from the meatus. His urethra would admit a No. 3, but it was exceedingly tight; and as he was desirous of returning to the country, I advised that the stricture should be immediately split. The dilator was at once passed, and the stricture split, so that a No. 12 catheter could be introduced into the bladder. The urine having been withdrawn, the catheter was removed, and he was directed to return home, and remain quiet the same afternoon. He again presented himself on the second day after the operation, stating that he had no constitutional disturbance, and that

the stream of urine was improved. A No. 12 catheter was again introduced, and without any drawback ; this was repeated on three occasions, when, as he had been previously in the habit of passing a No. 3, he had no difficulty in introducing a No. 12, and he returned to the country. I have had frequent opportunities of seeing this gentleman since, but he has never required to consult me again respecting his stricture.

Mr. H., a gentleman of colour, consulted me for stricture, situated at the membranous portion of the urethra. He had suffered from difficult micturition for about eighteen years ; there were, however, no complications, and his stricture would admit a No. 3, which for some years he had been in the habit of passing. As he was a nervous person, he was placed under the influence of chloroform, when the dilator was introduced, and the stricture split with the No. 12 tube. The No. 12 catheter was then introduced, and the urine removed. There was scarcely any bleeding ; he never had a bad symptom, or took medicine, and in a fortnight returned to Australia, passing his No. 12 catheter, which, with others, in case of fresh cause of difficulty, he procured from Messrs. Whicker and Blaize.

Mr. B., aged 65, residing at Hampstead, consulted me for stricture of thirty years' duration. He had been occasionally the subject of retention of urine, which was usually relieved by warm baths and opium. When I first saw him, in 1858, I could only introduce No. 1, and as he was in great dread of any operation, I consented to dilate the stricture in the ordinary manner. By the gradual process I eventually arrived at No. 8, when, from circumstances, he was compelled to discontinue his attendance; and upon his again consulting me, in the latter part of 1859, the stricture had so far returned that I was compelled to begin *de novo*. This time I persuaded him to have the stricture split, and, without the aid of chloroform the dilator was passed, and the stricture split to No. 13, his urethra being sufficiently capacious to take that size. Although an old man for his age, he never had a bad symptom, and ceased to continue under my care after my teaching him to pass his large-sized catheter, which he accomplished in about three weeks.

Is there, then, any special class of strictures where the operation by rupture is inadmissible, either from the situation, kind, and number of strictures, or from other complications? I be-

lieve not; but that there is a large number where any cutting operation must be attended with considerable danger, is now well known. Let us, for example, take a class of cases frequently met with, where fistulæ in perinæo, or elsewhere, exist, combined with great hypertrophy of all the tissues in the immediate contiguity of the stricture, and in which considerable time must be expended, and much skill employed before any instrument can be passed into the bladder. Here, to cut into a small groove requires great coolness, determination, and experience; and even when the operator is successful in reaching the groove, and so far dividing the obstruction as to admit a large catheter, the patient is confined to bed for many weeks before union will take place sufficiently to allow of the withdrawal of the instrument. And, again, what is the advantage of dividing a mass of hypertrophied tissue which does not encroach upon the canal, but necessarily intervenes between the knife and the urethra? If the diameter of the urethra can be enlarged to the same extent without such a proceeding, and a free outlet afforded for the urine, the hypertrophy will diminish, and the parts will eventually be restored to their original integrity.



But let me refer to another class where three or more strictures of varying density exist at different parts of the canal. Is the urethra to be laid open from end to end, or can the three obstructions be divided by internal incision at one and the same operation? I have seen these operations performed by the best surgeons in London, where no possible objection could be urged against their anatomical knowledge, their surgical experience, or their operative dexterity under the most trying circumstances; and I regret to say that they have not been without a fatal issue. No doubt it looks more surgical to place a patient in the position of lithotomy, and undertake a difficult piece of dissection in his perinæum, than to simply introduce an instrument, and pass a large tube, as it were, blindly between its blades, but the results of such cases warrant me in doubting the wisdom of such a proceeding, when the same result can be attained in another manner without difficulty or danger.

Hitherto I have only considered the applicability of the dilator, so far as rupturing or splitting the stricture is concerned, but it is equally efficacious where dilatation is desired, and possesses an advantage over every kind of bougie or sound in its power of dilating a stric-

ture to any required extent without being withdrawn. All practical surgeons know that having, for instance, had some difficulty in passing a No. 1, it is frequently impossible to introduce a No. 2 at the same visit, whereas, having once introduced the dilator, its diameter can be increased to any extent the surgeon may desire ; for this purpose it is not necessary that consecutive tubes should be passed, as by introducing at once a No. 8 or No. 9, and very gently pressing the tube onwards between the blades, they become separated to a considerable distance from the point of the tube, as is excellently shown in Figure 1 (page 4). By this means (which, however, occupies some little time) the stricture being acted upon from within, a greater amount of dilatation can be effected at one visit, and with much less pain to the patient than by passing consecutive bougies, which irritate the urethra, and owing to the abrupt increase of size, sometimes will not enter the stricture at all. If, in adopting the gradually dilating plan, the tube is very slowly passed, the pain will be trifling, and it will entirely cease, in the majority of instances, in a minute or two after the progress of the tube is discontinued. I cannot conceive any instrument more fitting, where gradual dilatation is

desired ; the blades can be expanded with the greatest nicety, in entire obedience to the desire of the surgeon, and the feelings of the patient, and as the dilatation is effected from within outwards, its force is expended in the most advantageous direction.

But, although dilatation may be employed where time is no object, and where the stricture is of a yielding character, it is utterly incapable of effecting what may be accomplished by passing the large tube at once, and splitting the stricture; the increased diameter being maintained by the after-passage of bougies. The rapidity with which the case will terminate, so far as only to require the passage of a bougie twice or thrice in the course of the year, will depend upon the number of strictures, and the irritability of the patient, more than it will upon their density. In some cases, although great force is required to split the obstruction, the urethra is to a certain extent insensible, and bears the after-passage of the bougie without the least manifestation of pain, whilst in others, more especially where there have been three or more impediments, and the urethra is very irritable, it is necessary to recede a size or two, owing to the spasms being so great during the introduc-

tion of the larger instrument, as to cause suffering.

I have lately had a marked example of this in an artilleryman, sent to me by my friend, Dr. Gallway, Surgeon-major of the Royal Artillery. The patient had been the subject of stricture about seven years; he had been through the Crimean war, and an inmate of the military hospital, more or less, ever since his return. Very many and long-continued attempts were made to get a catheter into his bladder without success, and Dr. Gallway requested me to meet him in consultation upon the case. I did so, but was equally unsuccessful in passing a catheter. I therefore requested Dr. Gallway to allow him to be admitted into the Westminster Hospital, to which he consented, and after several trials, I at last succeeded in passing No. 1 through four obstructions into the bladder. As the patient was excessively nervous and irritable, I delayed splitting the stricture until No. 2 was attained, when the stricture was split with the No. 10 tube, and a No. 10 catheter immediately passed: he did not suffer from the operation in the slightest degree, and his urine was passed in an improved stream. On the second day from the operation, I attempted to pass the No. 10, but the spasm was so per-

sistent that I preferred having recourse to No. 8 rather than give him pain; and I have been up to the present, now three weeks after the operation, content with that size, which passes with great ease, but still excites spasm. I have no question that in a little time No. 10 may be again used. The urethra must necessarily remain sore under such circumstances, and thus excites an amount of spasm which, by grasping the catheter, adds to the patient's suffering.

In conclusion I may add, that in advocating the treatment of stricture by rupture I claim simply that credit which attaches to the publication of a series of interesting cases (examples of many others) which have been subjected to this novel treatment. That the principle upon which the instrument is constructed is as old as the hills, and that the power of the wedge has been known as long as the simplest rules of mechanics have been taught, I freely admit, but I have yet to learn that that principle has been heretofore applied to the treatment of stricture of the urethra in the manner detailed above, and with such highly satisfactory results.

Shortly before the publication of this book, I had an opportunity of ascertaining what was the exact condition of the urethra when a stricture had been split after death.

A man was admitted under me into the Westminster Hospital in Oct., 1861. He had been the subject of stricture for eighteen years, and for the last twelve years the bladder had been relieved with increasing difficulty until he could only pass his water in drops. I endeavoured to introduce a No. 1 catheter, but found the stricture too contracted to admit even that size, and he was sent to bed, with directions to take castor-oil and remain quiet for a few days. In the meantime he was attacked with fever, and he died on Sunday, Nov. 10th. The friends objecting to a general *post-mortem* examination, I ultimately succeeded in persuading them to permit the urethra to be examined, which I was exceedingly anxious to do, for the purpose of ascertaining what was the precise effect produced by the passage of the dilator and its large tube. With some difficulty I introduced the dilator through two strictures, one within four inches of the meatus, and the second at the bulb, both apparently about half an inch long. The sensation was

like that I have frequently experienced during life, and the dilator having fairly reached the bladder, I passed the No. 12 tube, split the strictures, and before withdrawing it, rotated the dilator in the manner already described. My friend, Mr. Heath, now removed the penis, with a portion of the bladder and rectum, and having laid open the urethra, the two strictures, as is shown in the lithographic drawing, were found to be split at the under portion of the canal, the rent being directly in the median line, and limited to the extent of the obstruction. The contiguous structures were uninjured, and the divided mucous membrane gaped to an extent sufficient to permit the passage of a large bougie.

To Mr. Rawdon Macnamara I am indebted for the introduction of the dilator into the Meath Hospital; and to prove that success is not attributable to any facility I may possess in introducing the instrument and treating the cases, I will relate three most interesting instances that have come under that gentleman's professional care, and were published by him in a pamphlet entitled "On the Treatment of Stricture by the Immediate Plan." Mr. Macnamara, in his preface, says, "The more extended becomes my experience of this plan of treatment, the more convinced am I of its adaptability to almost every possible phase of this most troublesome disease, and the more satisfied of its great superiority over every other method heretofore suggested for its relief. In the following pages, I have dwelt on its simplicity, facility, security, and immunity from pain, hæmorrhage, and the varied sequelæ attendant on our other methods of treating stricture; here it but remains for me to repeat these assertions in a more emphatic manner. Every other day's experience of the "Immediate Plan" serves but still more to assure me that it is one of the greatest improvements in modern surgery."



In page 9, he says, "The first case on which the plan of forcible dilatation, so far as I am aware, was employed in this city, was that of M. E., aged 60. This man had been repeatedly under treatment for stricture of the urethra, and had been subjected to a false passage. He came into the Meath Hospital in the latter end of April, under Mr. Collis's care. After some days, we were able to introduce Mr. Holt's instrument; the stricture was burst on the 2nd of May, the patient went out on the 10th of the same month, and up to the present moment this individual is able to pass water in a full stream, and to admit of the introduction of a No. 10 catheter, and this without one single untoward symptom, from the period of the operation to that of his discharge."

The second case is one where a false passage had been previously made, and is thus recorded: "The case of A. B., aged 50, admitted into the Meath Hospital September 10th of the present year, was a most interesting one. He had long been subject to stricture, for the relief of which he had been treated by several surgeons on the plan of gradual dilatation. The largest sized instrument, however, that any of them ever succeeded in getting in, was

No. 4 ; and on the last occasion, in which an effort had been made to introduce an instrument, considerable difficulty was experienced, and a false passage was made, attended, as he states, with the loss of a considerable quantity of blood. After a great deal of trouble, I was enabled to get the dilator past the false passage into the bladder, split the stricture, introduce a No. 8 catheter, and empty the bladder ; the hæmorrhage was of the most trifling character, scarcely a drop of blood having been lost ; and as to the pain, he exhibited so little concern, that I thought he had not felt it, and it was only in reply to my question on the subject that he expressed himself as having felt what had been done to him. This operation was performed on the 12th of September ; that night he had rigors, but of a very slight nature. Next day, and ever since, he was as if he had undergone no operation, and I now experienced no difficulty in passing a No. 9 catheter ; and, indeed, so far as his stricture is concerned, he might have been discharged cured the day but one after Mr. Holt's instrument had been used. He still, however, remains in the Hospital, for the treatment of quite another disease, having no connexion whatsoever with the stricture."

The following case is one of great importance, both as to the gravity of the lesions, and in consequence of the manoeuvre employed, by which the dilator was introduced into the bladder.

R. T., aged 57, admitted into Meath Hospital in September, suffering from the effects of stricture of the urethra in three different situations, the most anterior being the tensest stricture it has ever been my lot to feel. This patient was under my friend Mr. Philip Crampton Smyly's care, to whose courtesy I am indebted for the liberty to make mention of it on the present occasion. This man had been on three different occasions operated on by different surgeons for the relief of his stricture, by external incision. He now is in a most deplorable state, Mr. Smyly with difficulty passing the smallest size instrument. On the 17th of September Mr. Smyly proceeded to put into operation the "Immediate Plan," and after prolonged manipulation, failed in getting the instrument more than through the most exterior of the three strictures, when its further progress was arrested, not by the distant strictures, but by the tension of the anterior stricture, which could be distinctly felt tightly pressing the instrument, and completely impeding its further

onward progress. Nor was I more fortunate in my endeavours to introduce it further, and the operation should have been abandoned, had we not, on consultation, agreed to burst this stricture, and then to proceed with the further steps of the operation, which Mr. Smyly accordingly carried into effect : he introduced the stilette, burst completely the anterior stricture, withdrew the stilette, closed again the blades of the dilator, continued its course successfully on into the bladder, again introduced the stilette, burst the other strictures, and on the withdrawal of the dilator introduced a No. 8 catheter into the bladder. In this case there was but little hæmorrhage, and but very trifling constitutional disturbance. This case made a profound impression on me at the time, as a good example of forcible dilatation *versus* external incision ; and the manœuvre by which the resistance of the anterior stricture was overcome, and the bladder at last reached, is one that merits attention, consideration, and recollection in similar cases at the hands of the operating surgeon."

To Mr. Smyly is the profession indebted for the first publication of the means of ascertaining with perfect precision when the dilator had entered the bladder. I, like that surgeon, had

frequently thought of how this could be accomplished ; but until lately had not devised a means so simple as not to complicate the facile performance of the operation. The published report of that gentleman, containing a drawing of his improvement is already before the profession ; and although I think it will be admitted that the instrument, as now altered by myself, embraces all the requirements in the most simple manner, yet I cannot but acknowledge the advantage that has arisen from Mr. Smyly's publication of his ingenious plan, which affords another instance of the readiness of our Irish brethren to advance the science and art of surgery.

London, New Burlington Street,  
March, 1867.

MESSRS. CHURCHILL & SONS'

Publications,

IN

M E D I C I N E

AND THE VARIOUS BRANCHES OF

NATURAL SCIENCE.



"It would be unjust to conclude this notice without saying a few words in favour of Mr. Churchill, from whom the profession is receiving, it may be truly said, the most beautiful series of Illustrated Medical Works which has ever been published."—*Lancet*.

"All the publications of Mr. Churchill are prepared with so much taste and neatness, that it is superfluous to speak of them in terms of commendation."—*Edinburgh Medical and Surgical Journal*.

"No one is more distinguished for the elegance and *recherché* style of his publications than Mr. Churchill."—*Provincial Medical Journal*.

"Mr. Churchill's publications are very handsomely got up: the engravings are remarkably well executed."—*Dublin Medical Press*.

"The typography, illustrations, and getting up are, in all Mr. Churchill's publications, most beautiful."—*Monthly Journal of Medical Science*.

"Mr. Churchill's illustrated works are among the best that emanate from the Medical Press."—*Medical Times*.

"We have before called the attention of both students and practitioners to the great advantage which Mr. Churchill has conferred on the profession, in the issue, at such a moderate cost, of works so highly creditable in point of artistic execution and scientific merit."—*Dublin Quarterly Journal*.



MESSRS. CHURCHILL & SONS are the Publishers of the following Periodicals, offering to Authors a wide extent of Literary Announcement, and a Medium of Advertisement, addressed to all Classes of the Profession. •

**THE BRITISH AND FOREIGN MEDICO-  
CHIRURGICAL REVIEW,  
AND  
QUARTERLY JOURNAL OF PRACTICAL MEDICINE  
AND SURGERY.**

Price Six Shillings. Nos. I. to LXXVII.

**THE QUARTERLY JOURNAL OF  
SCIENCE.**

Price Five Shillings. Nos. I. to XIII.

**THE QUARTERLY JOURNAL OF  
MICROSCOPICAL SCIENCE,**

INCLUDING THE TRANSACTIONS OF THE ROYAL  
MICROSCOPICAL SOCIETY OF LONDON.

Edited by DR. LANKESTER, F.R.S., and GEORGE  
BUSK, F.R.S. Price 4s. Nos. I. to XXV.

**THE JOURNAL OF MENTAL SCIENCE.**

By authority of the Medico-Psychological  
Association.

Edited by C. L. ROBERTSON, M.D., and HENRY  
MAUDSLEY, M.D.

Published Quarterly, price Half-a-Crown. *New  
Series.* Nos. I. to XXIV.

**ARCHIVES OF MEDICINE :**

A Record of Practical Observations and Anatomical and Chemical Researches, connected with the Investigation\* and Treatment of Disease. Edited by Dr. LIONEL S. BEALE, F.R.S. Published Quarterly; Nos. I. to VIII., 3s. 6d.; IX. to XII., 2s. 6d.; XIII. to XV., 3s.

**ARCHIVES OF DENTISTRY :**

Edited by EDWIN TRUMAN. Published Quarterly, price 4s. Nos. I. to IV.

**THE YEAR-BOOK OF PHARMACY  
AND  
CHEMISTS' DESK COMPANION  
FOR 1866.**

BEING A PRACTICAL SUMMARY OF RESEARCHES  
PHARMACY, MATERIA MEDICA,  
AND PHARMACEUTICAL CHEMISTRY, DURING  
THE YEAR 1865.

Edited by CHARLES H. WOOD, F.C.S., and  
CHAS. SHARP. Price 2s. 6d.

**THE ROYAL LONDON OPHTHALMIC HOSPITAL  
REPORTS, AND JOURNAL OF OPHTHALMIC  
MEDICINE AND SURGERY.**

Vol. V., Part 3,

**THE MEDICAL TIMES & GAZETTE.**

Published Weekly, price Sixpence, or Stamped,  
Sevenpence.

Annual Subscription, £1. 6s., or Stamped,  
£1. 10s. 4d., and regularly forwarded to all  
parts of the Kingdom.

**THE HALF-YEARLY ABSTRACT OF  
THE MEDICAL SCIENCES.**

Being a Digest of the Contents of the principal  
British and Continental Medical Works;  
together with a Critical Report of the Pro-  
gress of Medicine and the Collateral Sciences.  
Post 8vo. cloth, 6s. 6d. Vols. I. to XLIV.

**THE PHARMACEUTICAL JOURNAL,  
CONTAINING THE TRANSACTIONS OF THE PHAR-  
MACEUTICAL SOCIETY.**

Published Monthly, price One Shilling.

\* \* Vols. I. to XXV., bound in cloth, price  
12s. 6d. each.

**THE BRITISH JOURNAL OF DENTAL  
SCIENCE.**

Published Monthly, price One Shilling. Nos.  
I. to CXXVI.

**THE MEDICAL DIRECTORY FOR THE  
UNITED KINGDOM.**

Published Annually. 8vo. cloth, 10s. 6d.

**ANNALS OF MILITARY AND NAVAL  
SURGERY AND TROPICAL MEDICINE  
AND HYGIENE,**

Embracing the experience of the Medical Offi-  
cers of Her Majesty's Armies and Fleets in  
all parts of the World.

Vol. I., price 7s.

# A CLASSIFIED INDEX

TO

## MESSRS. CHURCHILL & SONS' CATALOGUE.

### ANATOMY.

	PAGE
Anatomical Remembrancer ..	7
Flower on Nerves .. ..	16
Hassall's Micros. Anatomy ..	19
Heale's Anatomy of the Lungs ..	19
Heath's Practical Anatomy ..	20
Hilden's Human Osteology ..	20
Do. on Dissections .. ..	20
Huxley's Comparative Anatomy ..	21
Jones' and Sieveking's Patho- logical Anatomy .. ..	22
MacDougal—Hirschfeld on the Nervous System .. ..	v
MacLise's Surgical Anatomy ..	25
St. Bartholomew's Hospital Catalogue .. ..	31
Sibson's Medical Anatomy ..	32
Waters' Anatomy of Lung ..	37
Wheeler's Anatomy for Artists ..	38
Wilson's Anatomy .. ..	39

### CHEMISTRY.

Bernays' Notes for Students ..	9
Bloxam's Chemistry .. ..	10
Bowman's Practical Chemistry ..	10
Do. Medical do. .. ..	10
Fowies' Manual of Chemistry ..	16
Do. Actonian Prize .. ..	16
Do. Qualitative Analysis ..	16
Frusenius' Chemical Analysis ..	16
Galloway's First Step .. ..	17
Do. Second Step .. ..	17
Do. Analysis .. ..	17
Do. Tables .. ..	17
Griffiths' Four Seasons .. ..	18
Horsley's Chem. Philosophy ..	21
Mulder on the Chemistry of Wine ..	27
Plattner & Muspratt on Blowpipe ..	28
Speer's Pathol. Chemistry ..	33
Sutton's Volumetric Analysis ..	34

### CLIMATE.

Aspinall on San Remo .. ..	7
Bennet's Winter in the South of Europe .. ..	9
Chambers on Italy .. ..	12
Dalrymple on Egypt .. ..	14
Francis on Change of Climate ..	16
Hall on Torquay .. ..	19
Haviland on Climate .. ..	19
Lee on Climate .. ..	24
Do. Watering Places of England ..	24
McClelland on Bengal .. ..	25
McNicol on Southport .. ..	25
Martin on Tropical Climates ..	26
Moore's Diseases of India ..	26
Scroesby-Jackson's Climatology ..	31
Shapter on South Devon .. ..	32
Sordet on Mentone .. ..	32
Taylor on Pau and Pyrenees ..	34

### DEFORMITIES, &c.

	PAGE
Adams on Spinal Curvature ..	6
Do. on Clubfoot .. ..	6
Biggs' Orthopraxy .. ..	10
Bishop on Deformities .. ..	10
Do. Articulate Sounds .. ..	10
Brodhurst on Spine .. ..	11
Do. on Clubfoot .. ..	11
Godfrey on Spine .. ..	17
Huguan on Hip Joint .. ..	21
Salt on Lower Extremities ..	31
Tamplin on Spine .. ..	34

### DISEASES OF WOMEN AND CHILDREN.

Ballard on Infants and Mothers ..	7
Bennet on Uterus .. ..	9
Bird on Children .. ..	10
Bryant's Surg. Diseases of Child ..	11
Eyre's Practical Remarks .. ..	15
Harrison on Children .. ..	19
Hood on Scarlet Fever, &c. ..	21
Kiwisch (ed. by Clay) on Ovaries ..	15
Lee's Ovarian & Uterine Diseases ..	24
Do. on Speculum .. ..	24
Ritchie on Ovaries .. ..	30
Seymour on Ovaria .. ..	32
Tilt on Uterine Inflammation ..	35
Do. Uterine Therapeutics .. ..	35
Do. on Change of Life .. ..	35
Underwood on Children .. ..	36
Wells on the Ovaries .. ..	33
West on Women .. ..	38
Do. (Uvedale) on Puerp. Diseases ..	38

### GENERATIVE ORGANS, Diseases of, and SYPHILIS.

Acton on Reproductive Organs ..	6
Coute on Syphilis .. ..	14
Gant on Bladder .. ..	17
Hutchinson on Inherited Syphilis ..	22
Judd on Syphilis .. ..	23
Lee on Syphilis .. ..	24
Parker on Syphilis .. ..	27
Wilson on Syphilis .. ..	39

### HYGIENE.

Armstrong on Naval Hygiene ..	7
Beale's Laws of Health .. ..	8
Do. Health and Disease .. ..	8

### HYGIENE—continued.

	PAGE
Carter on Training .. ..	12
Chayasse's Advice to a Mother ..	13
Do. Advice to a Wife .. ..	13
Dobell's Germs and Vestiges of Disease .. ..	15
Do. Diet and Regimen .. ..	15
File & Urquhart on Turkish Bath ..	16
Gordon on Army Hygiene .. ..	17
Granville on Vichy .. ..	18
Hartwig on Sea Bathing .. ..	19
Do. Physical Education .. ..	19
Hufeland's Art of prolonging Life ..	21
Lee's Baths of France, Germany, and Switzerland .. ..	24
Moore's Health in Tropics .. ..	26
Parkes on Hygiene .. ..	27
Parkin on Disease .. ..	28
Pearse's Notes on Health .. ..	29
Pickford on Hygiene .. ..	28
Robertson on Diet .. ..	30
Routh on Infant Feeding .. ..	30
Wells' Seamen's Medicine Chest ..	38
Wife's Domain .. ..	38
Wilson on Healthy Skin .. ..	39
Do. on Mineral Waters .. ..	39
Do. on Turkish Bath .. ..	39

### MATERIA MEDICA and PHARMACY.

Bateman's Magnacopia .. ..	8
Beasley's Formulary .. ..	9
Do. Receipt Book .. ..	9
Do. Book of Prescriptions ..	9
Frazer's Materia Medica .. ..	16
Nerins' Analysis of Pharmacop. ..	27
Pereira's Selecta & Prescripts ..	28
Prescriber's Pharmacopoeia .. ..	29
Royle's Materia Medica .. ..	31
Squire's Hospital Pharmacopoeias ..	33
Do. Companion to the Phar- macopoeia .. ..	33
Stegall's First Lines for Che- mists and Druggists .. ..	34
Stowe's Toxicological Chart .. ..	34
Taylor on Poisons .. ..	35
Waring's Therapeutics .. ..	37
Wittstein's Pharmacy .. ..	39

### MEDICINE.

Adams on Rheumatic Gout .. ..	6
Addison on Cell Therapeutics ..	6
Do. on Healthy and Dis- eased Structure .. ..	6
Aldis's Hospital Practice .. ..	6
Anderson (Andrew) on Fever ..	7
Do. (Thos.) on Yellow Fever ..	7



**MEDICINE—continued.**

	PAGE
Austin on Paralysis .. ..	7
Barclay on Medical Diagnosis ..	8
Do. on Gout .. ..	8
Barlow's Practice of Medicine ..	8
Basham on Dropsy .. ..	8
Brinton on Stomach .. ..	11
Do. on Intestinal Obstruction ..	11
Budd on the Liver .. ..	11
Do. on Stomach .. ..	11
Camplin on Diabetes .. ..	12
Chambers on the Indigestions ..	12
Do. Lectures .. ..	12
Cockle on Cancer .. ..	12
Davey's Ganglionic Nervous Syst.	15
Day's Clinical Histories .. ..	15
Eyre on Stomach .. ..	15
Foster on the Sphygmograph ..	16
Fuller on Rheumatism .. ..	16
Gairdner on Gout .. ..	17
Gibb on Throat .. ..	17
Do. on Laryngoscope .. ..	17
Graville on Sudden Death ..	18
Griffith on the Skin .. ..	18
Gully's Simple Treatment .. ..	18
Habershon on the Abdomen ..	18
Do. on Mercury .. ..	18
Hall (Marshall) on Apnoea ..	18
Do. Observations .. ..	18
Headland—Action of Medicines ..	19
Do. Medical Handbook .. ..	19
Hooper's Physician's Vade-Mecum .. ..	18
Inman's New Theory .. ..	22
Do. Myalgia .. ..	22
James on Laryngoscope .. ..	22
Jones (Bence) on Pathology and Therapeutics .. ..	22
MacLachlan on Advanced Life ..	25
MacLeod on Acolic Diseases ..	25
Marcel on Chronic Alcoholism ..	25
Macpherson on Cholera .. ..	26
Markham on Bleeding .. ..	26
Meryon on Paralysis .. ..	26
Musket on Apoplexy .. ..	27
Nicholson on Yellow Fever ..	27
Parkin on Cholera .. ..	28
Pavy on Diabetes .. ..	28
Pett's Principles and Practice of Medicine .. ..	28
Roberts on Palsy .. ..	30
Robertson on Gout .. ..	30
Sansom on Cholera .. ..	31
Savory's Compendium .. ..	31
Semple on Cough .. ..	32
Seymour on Dropsy .. ..	32
Shaw's Remembrancer .. ..	32
Shrimpton on Cholera .. ..	32
Smee on Debility .. ..	32
Thomas' Practice of Physic ..	35
Thudichum on Gall Stones ..	35
Todd's Clinical Lectures .. ..	36
Tweedie on Continued Fevers ..	36
Walker on Diphtheria .. ..	37
What to Observe at the Bedside ..	37
Williams' Principles .. ..	38
Wright on Headaches .. ..	39

**MICROSCOPE.**

Beale on Microscope in Medicine ..	8
Carpenter on Microscope .. ..	12
Schuch on do. .. ..	31

**MISCELLANEOUS.**

Acton on Prostitution .. ..	6
Barclay's Medical Errors .. ..	8

**MISCELLANEOUS—cont'd.**

	PAGE
Barker & Edwards' Photographs ..	8
Bascome on Epidemics .. ..	8
Bryce on Sebastopol .. ..	11
Buckle's Hospital Statistics ..	11
Cooley's Cyclopædia .. ..	13
Gordon on China .. ..	17
Graves' Physiology and Medicine ..	17
Guy's Hospital Reports .. ..	17
Harrison on Lead in Water ..	19
Hingeston's Topics of the Day ..	20
Howe on Epidemics .. ..	21
Lane's Hydropathy .. ..	23
Lee on Homœop. and Hydrop. ..	24
London Hospital Reports .. ..	25
Marcel on Food .. ..	25
Massy on Recruits .. ..	26
Mayne's Medical Vocabulary ..	26
Part's Case Book .. ..	28
Redwood's Supplement to Pharmacopæia .. ..	30
Ryan on Infanticide .. ..	31
St. George's Hospital Reports ..	31
Simms' Winter in Paris .. ..	32
Snow on Chloroform .. ..	33
Steggall's Medical Manual .. ..	34
Do. Gregory's Conspectus ..	34
Do. Celsus .. ..	34
Waring's Tropical Resident at Home .. ..	37
Whithead on Transmission ..	38

**NERVOUS DISORDERS AND INDIGESTION.**

Althaus on Epilepsy, Hysteria ..	7
Birch on Constipation .. ..	10
Carter on Hysteria .. ..	12
Downing on Neuralgia .. ..	15
Hunt on Heartburn .. ..	21
Jones (Handfield) on Functional Nervous Disorders .. ..	22
Leared on Imperfect Digestion ..	23
Lobb on Nervous Affections ..	24
Radcliffe on Epilepsy .. ..	29
Reynolds on the Brain .. ..	30
Do. on Epilepsy .. ..	30
Rowe on Nervous Diseases ..	31
Sieveling on Epilepsy .. ..	32
Turnbull on Stomach .. ..	36

**OBSTETRICS.**

Barnes on Placenta Prævia .. ..	8
Hodgeson on Puerperal Convulsions ..	20
Lee's Clinical Midwifery .. ..	24
Do. Consultations .. ..	24
Leishman's Mechanism of Parturition .. ..	24
Pretty's Aids during Labour ..	29
Priestley on Gravid Uterus ..	29
Ransbotham's Obstetrics .. ..	29
Do. Midwifery .. ..	30
Sinclair & Johnston's Midwifery ..	32
Smellie's Obstetric Plates .. ..	33
Smith's Manual of Obstetrics ..	33
Swayne's Aphorisms .. ..	34
Waller's Midwifery .. ..	37

**OPHTHALMOLOGY.**

Cooper on Injuries of Eye .. ..	13
Do. on Near Sight .. ..	13
Dalrymple on Eye .. ..	14

**OPHTHALMOLOGY—cont'd.**

	PAGE
Dixon on the Eye .. ..	15
Hogg on Ophthalmoscope .. ..	20
Hulke on the Ophthalmoscope ..	21
Jago on Entoptics .. ..	22
Jones' Ophthalmic Medicine ..	23
Do. Defects of Sight .. ..	23
Do. Eye and Ear .. ..	23
Macnamara on the Eye .. ..	25
Nunneley on the Organs of Vision ..	27
Solomon on Glaucoma .. ..	33
Walton on the Eye .. ..	37
Wells on Spectacles .. ..	37

**PHYSIOLOGY.**

Carpenter's Human .. ..	12
Do. Manual .. ..	12
Heale on Vital Causes .. ..	19
Richardson on Coagulation ..	30
Shea's Animal Physiology .. ..	32
Virchow's (ed. by Chance) Cellular Pathology .. ..	12

**PSYCHOLOGY.**

Arlidge on the State of Lunacy ..	7
Bucknill and Tuke's Psychological Medicine .. ..	12
Conolly on Asylums .. ..	13
Davey on Nature of Insanity ..	15
Dunn on Psychology .. ..	15
Hood on Criminal Lunatics ..	21
Millington on Treatment of Insane ..	26
Murray on Emotional Diseases ..	27
Noble on Mind .. ..	27
Sankey on Mental Diseases .. ..	31
Williams (J. H.) Unsoundness of Mind .. ..	38

**PULMONARY and CHEST DISEASES, &c.**

Alison on Pulmonary Consumption .. ..	6
Barker on the Lungs .. ..	8
Billing on Lungs and Heart ..	10
Bright on the Chest .. ..	11
Cotton on Consumption .. ..	14
Do. on Stethoscope .. ..	14
Davies on Lungs and Heart ..	15
Dobell on the Chest .. ..	15
Do. on Tuberculosis .. ..	15
Do. on Winter Cough .. ..	15
Fenwick on Consumption .. ..	16
Fuller on Chest .. ..	17
Do. on Heart .. ..	17
Jones (Jas.) on Consumption ..	22
Lacune on Auscultation .. ..	23
Markham on Heart .. ..	26
Pencock on the Heart .. ..	28
Richardson on Consumption ..	30
Salter on Asthma .. ..	31
Skoda on Auscultation .. ..	26
Thompson on Consumption ..	35
Timms on Consumption .. ..	35
Turnbull on Consumption .. ..	37
Waters on Emphysema .. ..	37
Weber on Auscultation .. ..	37

# CLASSIFIED INDEX.

## RENAL and URINARY

### DISEASES.

	PAGE
Acton on Urinary Organs ..	6
Beale on Urine .. ..	8
Bird's Urinary Deposits ..	10
Coulson on Bladder .. ..	14
Hassall on Urine .. ..	19
Parkes on Urine .. ..	27
Thudichum on Urine .. ..	35
Todd on Urinary Organs ..	36

### SCIENCE.

Baxter on Organic Polarity ..	8
Bentley's Manual of Botany ..	9
Bird's Natural Philosophy ..	10
Craig on Electric Tension ..	14
Hardwich's Photography ..	19
Hinds' Harmonies .. ..	20
Howard on the Clouds .. ..	21
Jones on Vision .. ..	23
Do. on Body, Sense, and Mind	23
Mayne's Lexicon .. ..	26
Noad on the Inductorium ..	27
Pratt's Genealogy of Creation ..	29
Do. Eccentric & Centric Force	29
Do. on Orbital Motion .. ..	29
Do. Astronomical Investigations	29
Do. Oracles of God .. ..	29
Rainey on Shells .. ..	29
Reymond's Animal Electricity	30
Taylor's Medical Jurisprudence	34
Unger's Botanical Letters ..	36
Vestiges of Creation .. ..	36

## SURGERY.

	PAGE
Adams on Reparation of Tendons	6
Do. Subcutaneous Surgery	6
Anderson on the Skin .. ..	7
Ashton on Rectum .. ..	7
Brodhurst on Anchylosis ..	11
Bryant on Diseases of Joints ..	11
Do. Clinical Surgery .. ..	11
Callender on Rupture .. ..	12
Chapman on Ulcers .. ..	12
Do. Varicose Veins .. ..	12
Clark's Outlines of Surgery ..	13
Collis on Cancer .. ..	13
Cooper (Sir A.) on Testis ..	14
Do. (S.) Surg. Dictionary	14
Coulson on Lithotomy .. ..	14
Curling on Rectum .. ..	14
Do. on Testis .. ..	14
Druitt's Surgeon's Vade-Mecum	15
Fayrer's Clinical Surgery ..	16
Fergusson's Surgery .. ..	16
Gamgee's Amputation at Hip-joint	17
Gant's Principles of Surgery ..	17
Heath's Minor Surgery and	
Bandaging .. ..	20
Higginbottom on Nitrate of Silver	20
Hodgson on Prostate .. ..	20
Holt on Stricture .. ..	21
James on Hernia .. ..	22
Jordan's Clinical Surgery ..	23
Lawrence's Surgery .. ..	23
Do. Ruptures .. ..	23
Lee on the Rectum, &c. .. ..	24
Liston's Surgery .. ..	24
Logan on Skin Diseases .. ..	24
Macleod's Surgical Diagnosis ..	25
Do. Surgery of the Crænea	25

## SURGERY—continued.

	PAGE
MacLise on Fractures .. ..	25
Maudsley's Operative Surgery ..	26
Naylor on Skin Diseases .. ..	27
Nunneley on Erysipelas .. ..	27
Pirrie's Surgery .. ..	28
Pirrie & Keith on Acupressure	28
Price on Excision of Knee-joint	29
Salt on Rupture .. ..	31
Sansom on Chloroform .. ..	31
Smith (Hy.) on Stricture .. ..	33
Do. on Hæmorrhoids .. ..	33
Do. on the Surgery of the Rectum	33
Do. (Dr. J.) Dental Anatomy	
and Surgery .. ..	33
Squire on Skin Diseases .. ..	33
Steggall's Surgical Manual ..	34
Teale on Amputation .. ..	35
Thompson on Stricture .. ..	35
Do. on Prostate .. ..	35
Do. Lithotomy and Lithotripsy	35
Tomes' Dental Surgery .. ..	36
Toynbee on Ear .. ..	36
Wade on Stricture .. ..	36
Webb's Surgeon's Ready Rules	37
Wilson on Skin Diseases .. ..	39
Do. Portraits of Skin Diseases	39
Yearsley on Deafness .. ..	39
Do. on Throat .. ..	39

## VETERINARY MEDICINE.

Blaine's Veterinary Art .. ..	10
Bourguignon on the Cattle Plague	10

TO BE COMPLETED IN TWELVE PARTS, 4TO., AT 7s. 6d. PER PART.

PARTS I. & II. NOW READY.

# A DESCRIPTIVE TREATISE ON THE NERVOUS SYSTEM OF MAN,

WITH THE MANNER OF DISSECTING IT.

By LUDOVIC HIRSCHFELD,

DOCTOR OF MEDICINE OF THE UNIVERSITIES OF PARIS AND WARSAW, PROFESSOR OF ANATOMY TO THE FACULTY OF MEDICINE OF WARSAW;

*Edited in English (from the French Edition)*

By ALEXANDER MASON MACDOUGAL, F.R.C.S.,

WITH

AN ATLAS OF ARTISTICALLY-COLOURED ILLUSTRATIONS,

Embracing the Anatomy of the entire Cerebro-Spinal and Sympathetic Nervous Centres and Distributions in their accurate relations with all the important Constituent Parts of the Human Economy, and embodied in a series of 56 Single and 9 Double Plates, comprising 197 Illustrations,

*Designed from Dissections prepared by the Author, and Drawn on Stone by*

J. B. LÉVEILLÉ.



MR. ACTON, M.R.C.S.

I.  
**A PRACTICAL TREATISE ON DISEASES OF THE URINARY AND GENERATIVE ORGANS IN BOTH SEXES.** Third Edition. 8vo. cloth, £1. 1s. With Plates, £1. 11s. 6d. The Plates alone, limp cloth, 10s. 6d.

II.  
**THE FUNCTIONS AND DISORDERS OF THE REPRODUCTIVE ORGANS IN CHILDHOOD, YOUTH, ADULT AGE, AND ADVANCED LIFE,** considered in their Physiological, Social, and Moral Relations. Fourth Edition. 8vo. cloth, 10s. 6d.

III.  
**PROSTITUTION:** Considered in its Moral, Social, and Sanitary Bearings, with a View to its Amelioration and Regulation. 8vo. cloth, 10s. 6d.

~~~~~  
DR. ADAMS, A.M.

**A TREATISE ON RHEUMATIC GOUT; OR, CHRONIC RHEUMATIC ARTHRITIS.** 8vo. cloth, with a Quarto Atlas of Plates, 21s.

MR. WILLIAM ADAMS, F.R.C.S.

I.  
**ON THE PATHOLOGY AND TREATMENT OF LATERAL AND OTHER FORMS OF CURVATURE OF THE SPINE.** With Plates. 8vo. cloth, 10s. 6d.

II.  
**CLUBFOOT:** its Causes, Pathology, and Treatment. Jacksonian Prize Essay for 1864. With 100 Engravings. 8vo. cloth, 12s.

III.  
**ON THE REPARATIVE PROCESS IN HUMAN TENDONS AFTER SUBCUTANEOUS DIVISION FOR THE CURE OF DEFORMITIES.** With Plates. 8vo. cloth, 6s.

IV.  
**SKETCH OF THE PRINCIPLES AND PRACTICE OF SUBCUTANEOUS SURGERY.** 8vo. cloth, 2s. 6d.

DR. WILLIAM ADDISON, F.R.S.

I.  
**CELL THERAPEUTICS.** 8vo. cloth, 4s.

II.  
**ON HEALTHY AND DISEASED STRUCTURE, AND THE TRUE PRINCIPLES OF TREATMENT FOR THE CURE OF DISEASE, ESPECIALLY CONSUMPTION AND SCROFULA,** founded on MICROSCOPICAL ANALYSIS. 8vo. cloth, 12s.

DR. ALDIS:

**AN INTRODUCTION TO HOSPITAL PRACTICE IN VARIOUS COMPLAINTS;** with Remarks on their Pathology and Treatment. 8vo. cloth, 5s. 6d.

DR. SOMERVILLE SCOTT ALISON, M.D. EDIN., F.R.C.P.

**THE PHYSICAL EXAMINATION OF THE CHEST IN PULMONARY CONSUMPTION, AND ITS INTERCURRENT DISEASES.** With Engravings. 8vo. cloth, 12s.

DR. ALTHAUS, M.D., M.R.C.P.

ON EPILEPSY, HYSTERIA, AND ATAXY. Cr. 8vo. cloth, 4s.

THE ANATOMICAL REMEMBRANCER; OR, COMPLETE  
POCKET ANATOMIST. Sixth Edition, carefully Revised. 32mo. cloth, 3s. 6d.

DR. MCCALL ANDERSON, M.D.

I.  
PARASITIC AFFECTIONS OF THE SKIN. With Engravings.  
8vo. cloth, 5s.

II.  
ECZEMA. 8vo. cloth, 5s.

III.  
PSORIASIS AND LEPRO. With Chromo-lithograph. 8vo. cloth, 5s.

DR. ANDREW ANDERSON, M.D.

TEN LECTURES INTRODUCTORY TO THE STUDY OF FEVER.  
Post 8vo. cloth, 5s.

DR. THOMAS ANDERSON, M.D.

HANDBOOK FOR YELLOW FEVER: ITS PATHOLOGY AND  
TREATMENT. To which is added a brief History of Cholera, and a method of Cure.  
Fcap. 8vo. cloth, 3s.

DR. ARLIDGE.

ON THE STATE OF LUNACY AND THE LEGAL PROVISION  
FOR THE INSANE; with Observations on the Construction and Organisation of  
Asylums. 8vo. cloth, 7s.

DR. ALEXANDER ARMSTRONG, R.N.

OBSERVATIONS ON NAVAL HYGIENE AND SCURVY.  
More particularly as the latter appeared during a Polar Voyage. 8vo. cloth, 5s.

MR. T. J. ASHTON.

I.  
ON THE DISEASES, INJURIES, AND MALFORMATIONS  
OF THE RECTUM AND ANUS. Fourth Edition. 8vo. cloth, 8s.

II.  
PROLAPSUS, FISTULA IN ANO, AND HÆMORRHOIDAL  
AFFECTIONS; their Pathology and Treatment. Second Edition. Post 8vo. cloth, 2s. 6d.

MR. W. B. ASPINALL.

SAN REMO AS A WINTER RESIDENCE. With Coloured Plates.  
Foolscap 8vo. cloth, 4s. 6d.

MR. THOS. J. AUSTIN, M.R.C.S.ENG.

A PRACTICAL ACCOUNT OF GENERAL PARALYSIS:  
Its Mental and Physical Symptoms, Statistics, Causes, Seat, and Treatment. 8vo. cloth, 6s.

DR. THOMAS BALLARD, M.D.

A NEW AND RATIONAL EXPLANATION OF THE DIS-  
EASES PECULIAR TO INFANTS AND MOTHERS; with obvious Suggestions  
for their Prevention and Cure. Post 8vo. cloth, 4s. 6d.

DR. BARCLAY.

I.  
A MANUAL OF MEDICAL DIAGNOSIS. Second Edition.  
Folscap 8vo. cloth, 8s. 6d.

II.  
MEDICAL ERRORS.—Fallacies connected with the Application of the  
Inductive Method of Reasoning to the Science of Medicine. Post 8vo. cloth, 5s.

III.  
GOUT AND RHEUMATISM IN RELATION TO DISEASE  
OF THE HEART. Post 8vo. cloth, 5s.

~~~~~  
DR. T. HERBERT BARKER, M.D., F.R.S., & MR. ERNEST EDWARDS, B.A.  
PHOTOGRAPHS OF EMINENT MEDICAL MEN, with brief  
Analytical Notices of their Works. Nos. I. to VIII., price 3s. each.

DR. W. G. BARKER, M.D.LOND.

ON DISEASES OF THE RESPIRATORY PASSAGES AND  
LUNGS, SPORADIC AND EPIDEMIC; their *Causes*, Pathology, Symptoms,  
and Treatment. Crown 8vo. cloth, 6s.

DR. BARLOW.

A MANUAL OF THE PRACTICE OF MEDICINE. Second  
Edition. Fcap. 8vo. cloth, 12s. 6d.

DR. BARNES.

THE PHYSIOLOGY AND TREATMENT OF PLACENTA  
PRÆVIA; being the Lettsonian Lectures on Midwifery for 1857. Post 8vo. cloth, 6s.

~~~~~  
DR. BASCOMB.

† A HISTORY OF EPIDEMIC PESTILENCES, FROM THE  
EARLIEST AGES. 8vo. cloth, 8s. †

DR. BASHAM.

ON DROPSY, AND ITS CONNECTION WITH DISEASES OF  
THE KIDNEYS, HEART, LUNGS AND LIVER. With 16 Plates. Third  
Edition. 8vo. cloth, 12s. 6d.

~~~~~  
MR. H. F. BAXTER, M.R.C.S.L.

ON ORGANIC POLARITY; showing a Connexion to exist between  
Organic Forces and Ordinary Polar Forces. Crown 8vo. cloth, 5s.

~~~~~  
MR. BATEMAN.

MAGNACOPIA: A Practical Library of Profitable Knowledge, commu-  
nicating the general Minutiæ of Chemical and Pharmaceutic Routine, together with the  
generality of Secret Forms of Preparations. Third Edition. 18mo. 6s.

MR. LIONEL J. BEALE, M.R.C.S.

I.  
THE LAWS OF HEALTH IN THEIR RELATIONS TO MIND  
AND BODY. A Series of Letters from an Old Practitioner to a Patient. Post 8vo.  
cloth, 7s. 6d.

II.  
HEALTH AND DISEASE, IN CONNECTION WITH THE  
GENERAL PRINCIPLES OF HYGIENE. Fcap. 8vo., 2s. 6d.

MESSRS. CHURCHILL & SONS' PUBLICATIONS.

DR. BEALE, F.R.S.

I.

URINE, URINARY DEPOSITS, AND CALCULI: and on the Treatment of Urinary Diseases. Numerous Engravings. Second Edition, much Enlarged. Post 8vo. cloth, 8s. 6d.

II.

THE MICROSCOPE, IN ITS APPLICATION TO PRACTICAL MEDICINE. Third Edition. With 58 Plates. 8vo. cloth, 16s.

III.

ILLUSTRATIONS OF THE SALTS OF URINE, URINARY DEPOSITS, and CALCULI. 37 Plates, containing upwards of 170 Figures copied from Nature, with descriptive Letterpress. 8vo. cloth, 9s. 6d.

MR. BEASLEY.

I.

THE BOOK OF PRESCRIPTIONS; containing 3000 Prescriptions. Collected from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Third Edition. 18mo. cloth, 6s.

II.

THE DRUGGIST'S GENERAL RECEIPT-BOOK; comprising a copious Veterinary Formulary and Table of Veterinary Materia Medica; Patent and Proprietary Medicines, Druggists' Nostrums, &c.; Perfumery, Skin Cosmetics, Hair Cosmetics, and Teeth Cosmetics; Beverages, Dietetic Articles, and Condiments; Trade Chemicals, Miscellaneous Preparations and Compounds used in the Arts, &c.; with useful Memoranda and Tables. Sixth Edition. 18mo. cloth, 6s.

III.

THE POCKET FORMULARY AND SYNOPSIS OF THE BRITISH AND FOREIGN PHARMACOPŒIAS; comprising standard and approved Formulæ for the Preparations and Compounds employed in Medical Practice. Eighth Edition, corrected and enlarged. 18mo. cloth, 6s.

DR. HENRY BENNET.

I.

A PRACTICAL TREATISE ON. INFLAMMATION AND OTHER DISEASES OF THE UTERUS. Fourth Edition, revised, with Additions. 8vo. cloth, 16s.

II.

A REVIEW OF THE PRESENT STATE (1856) OF UTERINE PATHOLOGY. 8vo. cloth, 4s.

III.

WINTER IN THE SOUTH OF EUROPE; OR, MENTONE, THE RIVIERA, CORSICA, SICILY, AND BIARRITZ, AS WINTER CLIMATES. Third Edition, with numerous Plates, Maps, and Wood Engravings. Post 8vo. cloth, 10s. 6d.

PROFESSOR BENTLEY, F.L.S.

A MANUAL OF BOTANY. With nearly 1,200 Engravings on Wood. Fcap. 8vo. cloth, 12s. 6d.

DR. BERNAYS.

NOTES FOR STUDENTS IN CHEMISTRY; being a Syllabus compiled from the Manuals of Miller, Fowries, Berzelius, Gerhardt, Goup-Besanez, &c. Fourth Edition. Fcap. 8vo. cloth, 3s.

MR. HENRY HEATHER BIGG.

† **ORTHOPRAXY:** the Mechanical Treatment of Deformities, Debilities, and Deficiencies of the Human Frame. With Engravings. Post 8vo. cloth, 10s.

DR. BILLING, F.R.S.

**ON DISEASES OF THE LUNGS AND HEART.** 8vo. cloth, 6s.

DR. S. B. BIRCH, M.D.

**CONSTIPATED BOWELS:** the Various Causes and the Rational Means of Cure. Second Edition. Post 8vo. cloth, 3s. 6d.

DR. GOLDING BIRD, F.R.S.

I.  
**URINARY DEPOSITS; THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS.** With Engravings. Fifth Edition. Edited by E. LLOYD BIRKETT, M.D. Post 8vo. cloth, 10s. 6d.

II.  
**ELEMENTS OF NATURAL PHILOSOPHY;** being an Experimental Introduction to the Study of the Physical Sciences. With numerous Engravings. Fifth Edition. Edited by CHARLES BROOKE, M.B. Cantab., F.R.S. Fcap. 8vo. cloth, 12s. 6d.

MR. BISHOP, F.R.S.

I.  
**ON DEFORMITIES OF THE HUMAN BODY,** their Pathology and Treatment. With Engravings on Wood. 8vo. cloth, 10s.

II.  
**ON ARTICULATE SOUNDS, AND ON THE CAUSES AND CURE OF IMPEDIMENTS OF SPEECH.** 8vo. cloth, 4s.

MR. P. HINCKES BIRD, F.R.C.S.

**PRACTICAL TREATISE ON THE DISEASES OF CHILDREN AND INFANTS AT THE BREAST.** Translated from the French of M. BOUCHUT, with Notes and Additions. 8vo. cloth, 20s.

MR. BLAINE.

**OUTLINES OF THE VETERINARY ART; OR, A TREATISE ON THE ANATOMY, PHYSIOLOGY, AND DISEASES OF THE HORSE, NEAT CATTLE, AND SHEEP.** Seventh Edition. By Charles Steel, M.R.C.V.S.L. With Plates. 8vo. cloth, 18s.

MR. BLOXAM.

**CHEMISTRY, INORGANIC AND ORGANIC;** with Experiments and a Comparison of Equivalent and Molecular Formulæ. With 276 Engravings on Wood. 8vo. cloth, 16s.

DR. BOURGUIGNON.

**ON THE CATTLE PLAGUE; OR, CONTAGIOUS TYPHUS IN HORNED CATTLE:** its History, Origin, Description, and Treatment. Post 8vo. 5s.

MR. JOHN E. BOWMAN, & MR. C. L. BLOXAM.

I.  
**PRACTICAL CHEMISTRY,** including Analysis. With numerous Illustrations on Wood. Fifth Edition. Foolscap 8vo. cloth, 6s. 6d.

II.  
**MEDICAL CHEMISTRY;** with Illustrations on Wood. Fourth Edition, carefully revised. Fcap. 8vo. cloth, 6s. 6d.

DR. JAMES BRIGHT.

ON DISEASES OF THE HEART, LUNGS, & AIR PASSAGES;  
with a Review of the several Climates recommended in these Affections. Third Edition. Post 8vo. cloth, 9s.

---

DR. BRINTON, F.R.S.

I.  
THE DISEASES OF THE STOMACH, with an Introduction on its  
Anatomy and Physiology; being Lectures delivered at St. Thomas's Hospital. Second  
Edition. 8vo. cloth, 10s. 6d.

II.

INTESTINAL OBSTRUCTION. Edited by DR. BUZZARD. Post 8vo.  
cloth, 5s.

---

MR. BERNARD E. BRODHURST, F.R.C.S.

I.  
CURVATURES OF THE SPINE: their Causes, Symptoms, Pathology,  
and Treatment. Second Edition. Roy. 8vo. cloth, with Engravings, 7s. 6d.

II.

ON THE NATURE AND TREATMENT OF CLUBFOOT AND  
ANALOGOUS DISTORTIONS involving the TIBIO-TARSAL ARTICULATION.  
With Engravings on Wood. 8vo. cloth, 4s. 6d.

III.

PRACTICAL OBSERVATIONS ON THE DISEASES OF THE  
JOINTS INVOLVING ANCHYLOSIS, and on the TREATMENT for the  
RESTORATION of MOTION. Third Edition, much enlarged, 8vo. cloth, 4s. 6d.

MR. THOMAS BRYANT, F.R.C.S.

I.  
ON THE DISEASES AND INJURIES OF THE JOINTS.  
CLINICAL AND PATHOLOGICAL OBSERVATIONS. Post 8vo. cloth, 7s. 6d.

II.

THE SURGICAL DISEASES OF CHILDREN. The Lettsomian  
Lectures, delivered March, 1863. Post 8vo. cloth, 5s.

III.

CLINICAL SURGERY. Parts I. to VII. 8vo., 3s. 6d. each.

---

DR. BRYCE.

ENGLAND AND FRANCE BEFORE SEBASTOPOL, looked at  
from a Medical Point of View. 8vo. cloth, 6s.

DR. BUCKLE, M.D., L.R.C.P. LOND.

VITAL AND ECONOMICAL STATISTICS OF THE HOSPITALS,  
INFIRMARIES, &c., OF ENGLAND AND WALES. Royal 8vo. 5s.

---

DR. BUDD, F.R.S.

I.  
ON DISEASES OF THE LIVER.

Illustrated with Coloured Plates and Engravings on Wood. Third Edition. 8vo. cloth, 16s.

II.

ON THE ORGANIC DISEASES AND FUNCTIONAL DIS-  
ORDERS OF THE STOMACH. 8vo. cloth, 9s.



MESSRS. CHURCHILL & SONS' PUBLICATIONS.

DR. JOHN CHARLES BUCKNILL, F.R.S., & DR. DANIEL H. TUKE.

**A MANUAL OF PSYCHOLOGICAL MEDICINE:** containing the History, Nosology, Description, Statistics, Diagnosis, Pathology, and Treatment of Insanity. Second Edition. 8vo. cloth, 15s.

MR. CALLENDER, F.R.C.S.

**FEMORAL RUPTURE:** Anatomy of the Parts concerned. With Plates. 8vo. cloth, 4s.

DR. JOHN M. CAMPLIN, F.L.S.

**ON DIABETES, AND ITS SUCCESSFUL TREATMENT.** Third Edition, by Dr. Glover. Fcap. 8vo. cloth, 3s. 6d.

MR. ROBERT B. CARTER, M.R.C.S.

**ON THE INFLUENCE OF EDUCATION AND TRAINING**  
IN PREVENTING DISEASES OF THE NERVOUS SYSTEM. Fcap. 8vo., 6s.

**THE PATHOLOGY AND TREATMENT OF HYSTERIA.** Post 8vo. cloth, 4s. 6d.

DR. CARPENTER, F.R.S.

**PRINCIPLES OF HUMAN PHYSIOLOGY.** With numerous Illustrations on Steel and Wood. Sixth Edition. Edited by Mr. HENRY POWER. 8vo. cloth, 26s.

**A MANUAL OF PHYSIOLOGY.** With 252 Illustrations on Steel and Wood. Fourth Edition. Fcap. 8vo. cloth, 12s. 6d.

**THE MICROSCOPE AND ITS REVELATIONS.** With numerous Engravings on Steel and Wood. Third Edition. Fcap. 8vo. cloth, 12s. 6d.

DR. CHAMBERS.

**LECTURES, CHIEFLY CLINICAL.** Fourth Edition. 8vo. cloth, 14s.

**THE INDIGESTIONS OR DISEASES OF THE DIGESTIVE**  
ORGANS FUNCTIONALLY TREATED. 8vo. cloth, 10s.

**SOME OF THE EFFECTS OF THE CLIMATE OF ITALY.** Crown 8vo. cloth, 4s. 6d.

DR. CHANCE, M.B.

**VIRCHOW'S CELLULAR PATHOLOGY, AS BASED UPON**  
PHYSIOLOGICAL AND PATHOLOGICAL HISTOLOGY. With 144 Engravings on Wood. 8vo. cloth, 16s.

MR. H. T. CHAPMAN, F.R.C.S.

**THE TREATMENT OF OBSTINATE ULCERS AND CUTA-**  
NEOUS ERUPTIONS OF THE LEG WITHOUT CONFINEMENT. Third Edition. Post 8vo. cloth, 3s. 6d.

II.

**VARICOSE VEINS:** their Nature, Consequences, and Treatment, Palliative and Curative. Second Edition. Post 8vo. cloth, 3s. 6d.

MESSRS. CHURCHILL & SONS' PUBLICATIONS.

MR. PYE HENRY CHAVASSE, F.R.C.S.

I.  
ADVICE TO A MOTHER ON THE MANAGEMENT OF  
HER CHILDREN. Eighth Edition. Foolscap 8vo., 2s. 6d.

II.  
ADVICE TO A WIFE ON THE MANAGEMENT OF HER  
OWN HEALTH. With an Introductory Chapter, especially addressed to a Young  
Wife. Seventh Edition. Fcap. 8vo., 2s. 6d.

MR. LE GROS CLARK, F.R.C.S.

OUTLINES OF SURGERY ; being an Epitome of the Lectures on the  
Principles and the Practice of Surgery, delivered at St. Thomas's Hospital. Fcap. 8vo.  
cloth, 5s.

MR. JOHN CLAY, M.R.C.S.

KIWISCH ON DISEASES OF THE OVARIES: Translated, by  
permission, from the last German Edition of his Clinical Lectures on the Special Patho-  
logy and Treatment of the Diseases of Women. With Notes, and an Appendix on the  
Operation of Ovariectomy. Royal 12mo. cloth, 16s.

DR. COCKLE, M.D.

ON INTRA-THORACIC CANCER. 8vo. 6s. 6d.

MR. COLLIS, M.B.DUB., F.R.C.S.I.

THE DIAGNOSIS AND TREATMENT OF CANCER AND  
THE TUMOURS ANALOGOUS TO IT. With coloured Plates. 8vo. cloth, 14s.

• DR. CONOLLY.

THE CONSTRUCTION AND GOVERNMENT OF LUNATIC  
ASYLUMS AND HOSPITALS FOR THE INSANE. With Plans. Post 8vo.  
cloth, 6s.

MR. COOLEY.

COMPREHENSIVE SUPPLEMENT TO THE PHARMACOPEIAS.

THE CYCLOPÆDIA OF PRACTICAL RECEIPTS, PRO-  
CESSES, AND COLLATERAL INFORMATION IN THE ARTS, MANU-  
FACTURES, PROFESSIONS, AND TRADES, INCLUDING MEDICINE,  
PHARMACY, AND DOMESTIC ECONOMY ; designed as a General Book of  
Reference for the Manufacturer, Tradesman, Amateur, and Heads of Families. Fourth  
and greatly enlarged Edition, 8vo. cloth, 28s.

MR. W. WHITE COOPER.

I.  
ON WOUNDS AND INJURIES OF THE EYE. Illustrated by  
17 Coloured Figures and 41 Woodcuts. 8vo. cloth, 12s.

II.  
ON NEAR SIGHT, AGED SIGHT, IMPAIRED VISION,  
AND THE MEANS OF ASSISTING SIGHT. With 31 Illustrations on Wood.  
• Second Edition. Fcap. 8vo. cloth, 7s. 6d.

SIR ASTLEY COOPER, BART., F.R.S.  
**ON THE STRUCTURE AND DISEASES OF THE TESTIS.**  
 With 24 Plates. Second Edition. Royal 4to., 20s.

~~~~~  
 MR. COOPER.

**A DICTIONARY OF PRACTICAL SURGERY AND ENCYCLO-  
 PÆDIA OF SURGICAL SCIENCE.** New Edition, brought down to the present  
 time. By SAMUEL A. LANE, F.R.C.S., assisted by various eminent Surgeons. Vol. I.,  
 8vo. cloth, £1. 5s.

~~~~~  
 MR. HOLMES COOTE, F.R.C.S.

**A REPORT ON SOME IMPORTANT POINTS IN THE  
 TREATMENT OF SYPHILIS.** 8vo. cloth, 5s.

DR. COTTON.

I.  
**ON CONSUMPTION:** Its Nature, Symptoms, and Treatment. To  
 which Essay was awarded the Fothergillian Gold Medal of the Medical Society of  
 London. Second Edition. 8vo. cloth, 8s.

II.  
**PHTHISIS AND THE STETHOSCOPE; OR, THE PHYSICAL  
 SIGNS OF CONSUMPTION.** Third Edition. Foolsap 8vo. cloth, 3s.

MR. COULSON.

I.  
**ON DISEASES OF THE BLADDER AND PROSTATE GLAND.**  
 New Edition, revised. *In Preparation.*

II.  
**ON LITHOTRITY AND LITHOTOMY;** with Engravings on Wood.  
 8vo. cloth, 8s.

~~~~~  
 MR. WILLIAM CRAIG, L.F.P.S., GLASGOW.

**ON THE INFLUENCE OF VARIATIONS OF ELECTRIC  
 TENSION AS THE REMOTE CAUSE OF EPIDEMIC AND OTHER  
 DISEASES.** 8vo. cloth, 10s.

~~~~~  
 MR. CURLING, F.R.S.

I.  
**OBSERVATIONS ON DISEASES OF THE RECTUM.** Third  
 Edition. 8vo. cloth, 7s. 6d.

II.  
**A PRACTICAL TREATISE ON DISEASES OF THE TESTIS,  
 SPERMATIC CORD, AND SCROTUM.** Third Edition, with Engravings. 8vo.  
 cloth, 16s.

~~~~~  
 DR. DALRYMPLE, M.R.C.P., F.R.C.S.

**THE CLIMATE OF EGYPT: METEOROLOGICAL AND MEDI-  
 CAL OBSERVATIONS,** with Practical Hints for Invalid Travellers. Post 8vo. cloth, 4s.

~~~~~  
 MR. JOHN DALRYMPLE, F.R.S., F.R.C.S.

**PATHOLOGY OF THE HUMAN EYE.** Complete in Nine Fasciculi:  
 imperial 4to., 20s. each; half-bound morocco, gilt tops, 9l. 15s.

DR. HERBERT DAVIES.

ON THE PHYSICAL DIAGNOSIS OF DISEASES OF THE LUNGS AND HEART. Second Edition. Post 8vo. cloth, 8s. †

DR. DAVEY.

I.  
THE GANGLIONIC NERVOUS SYSTEM: its Structure, Functions, and Diseases. 8vo. cloth, 9s.

II.  
ON THE NATURE AND PROXIMATE CAUSE OF SANITY.

DR. HENRY DAY, M.D., M.R.C.P.

CLINICAL HISTORIES; with Comments. 8vo. cloth, 7s. 6d.

MR. DIXON.

A GUIDE TO THE PRACTICAL STUDY OF DISEASES OF THE EYE. Third Edition. Post 8vo. cloth, 9s.

DR. DOBELL.

I.  
DEMONSTRATIONS OF DISEASES IN THE CHEST, AND THEIR PHYSICAL DIAGNOSIS. With Coloured Plates. 8vo. cloth, 12s. 6d. †

II.  
LECTURES ON THE GERMS AND VESTIGES OF DISEASE, and on the Prevention of the Invasion and Fatality of Disease by Periodical Examinations. 8vo. cloth, 6s. 6d.

III.  
A MANUAL OF DIET AND REGIMEN FOR PHYSICIAN AND PATIENT. Third Edition (for the year 1865). Crown 8vo. cloth, 1s. 6d.

IV.  
ON TUBERCULOSIS: ITS NATURE, CAUSE, AND TREATMENT; with Notes on Pancreatic Juice. Second Edition. Crown 8vo. cloth, 3s. 6d. x

V.  
ON WINTER COUGH (CATARRH, BRONCHITIS, EMPHYSEMA, ASTHMA); with an Appendix on some Principles of Diet in Disease—Lectures delivered at the Royal Infirmary for Diseases of the Chest. Post 8vo. cloth,

DR. TOOGOOD DOWNING.

NEURALGIA: its various Forms, Pathology, and Treatment. THE JACKSONIAN PRIZE ESSAY FOR 1850. 8vo. cloth, 10s. 6d.

DR. DRUITT, F.R.C.S.

THE SURGEON'S VADE-MECUM; with numerous Engravings on Wood. Ninth Edition. Foolscap 8vo. cloth, 12s. 6d.

MR. DUNN, F.R.C.S.

PSYCHOLOGY—PHYSIOLOGICAL, 4s.; MEDICAL, 3s.

SIR JAMES EYRE, M.D.

I.  
THE STOMACH AND ITS DIFFICULTIES. Fifth Edition. Fcap. 8vo. cloth, 2s. 6d.

II.  
PRACTICAL REMARKS ON SOME EXHAUSTING DISEASES. Second Edition. Post 8vo. cloth, 4s. 6d.

DR. FAYRER, M.D., F.R.C.S.

CLINICAL SURGERY IN INDIA. With Engravings. 8vo. cloth, 16s.

DR. FENWICK.

ON SCROFULA AND CONSUMPTION. Clergyman's Sore Throat, Catarrh, Croup, Bronchitis, Asthma. Fcap. 8vo., 2s. 6d.

SIR WILLIAM FERGUSSON, BART., F.R.S.

A SYSTEM OF PRACTICAL SURGERY; with numerous Illustrations on Wood. Fourth Edition. Fcap. 8vo. cloth, 12s. 6d.

SIR JOHN FIFE, F.R.C.S. AND MR. URQUHART.

MANUAL OF THE TURKISH BATH. Heat a Mode of Cure and a Source of Strength for Men and Animals. With Engravings. Post 8vo. cloth, 5s.

MR. FLOWER, F.R.S., F.R.C.S.

DIAGRAMS OF THE NERVES OF THE HUMAN BODY, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. Folio, containing Six Plates, 14s.

DR. BALTHAZAR FOSTER, M.D., M.R.C.P.

THE USE OF THE SPIGMOGRAPH IN THE INVESTIGATION OF DISEASE. With Engravings. 8vo. cloth, 2s. 6d.

MR. FOWNES, PH.D., F.R.S.

A MANUAL OF CHEMISTRY; with 187 Illustrations on Wood. Ninth Edition. Fcap. 8vo. cloth, 12s. 6d.

Edited by H. BENGE JONES, M.D., F.R.S., and A. W. HOFMANN, PH.D., F.R.S.

CHEMISTRY, AS EXEMPLIFYING THE WISDOM AND BENEFICENCE OF GOD. Second Edition. Fcap. 8vo. cloth, 4s. 6d.

INTRODUCTION TO QUALITATIVE ANALYSIS. Post 8vo. cloth, 2s.

DR. D. J. T. FRANCIS.

CHANGE OF CLIMATE; considered as a Remedy in Dyspeptic, Pulmonary, and other Chronic Affections; with an Account of the most Eligible Places of Residence for Invalids, at different Seasons of the Year. Post 8vo. cloth, 6s. 6d.

DR. W. FRAZER.

ELEMENTS OF MATERIA MEDICA; containing the Chemistry and Natural History of Drugs—their Effects, Doses, and Adulterations. Second Edition. 8vo. cloth, 10s. 6d.

C. REMIGIUS FRESENIUS.

A SYSTEM OF INSTRUCTION IN CHEMICAL ANALYSIS, Edited by LLOYD BULLOCK, F.C.S.

QUALITATIVE. Sixth Edition, with Coloured Plate illustrating Spectrum Analysis. 8vo. cloth, 10s. 6d.—QUANTITATIVE. Fourth Edition. 8vo. cloth, 10s.

DR. FULLER.

I.

ON DISEASES OF THE CHEST, including Diseases of the Heart and Great Vessels. With Engravings. 8vo. cloth, 12s. 6d.

II.

ON DISEASES OF THE HEART AND GREAT VESSELS. 8vo. cloth, 7s. 6d.

III.

ON RHEUMATISM, RHEUMATIC GOUT, AND SCIATICA: their Pathology, Symptoms, and Treatment. Third Edition. 8vo. cloth, 12s. 6d.

MR. GALLOWAY.

I.

THE FIRST STEP IN CHEMISTRY. Third Edition. Fcap. 8vo. cloth, 5s.

II.

THE SECOND STEP IN CHEMISTRY; or, the Student's Guide to the Higher Branches of the Science. With Engravings. 8vo. cloth, 10s.

III.

A MANUAL OF QUALITATIVE ANALYSIS. Fourth Edition. Post 8vo. cloth, 6s. 6d.

IV.

CHEMICAL TABLES. On Five Large Sheets, for School and Lecture Rooms. Second Edition. 4s. 6d.

MR. J. SAMPSON GAMGEE.

HISTORY OF A SUCCESSFUL CASE OF AMPUTATION AT THE HIP-JOINT (the limb 48-in. in circumference, 99 pounds weight). With 4 Photographs. 4to cloth, 10s. 6d.

MR. F. J. GANT, F.R.C.S.

I.

THE PRINCIPLES OF SURGERY: Clinical, Medical, and Operative. With Engravings. 8vo. cloth, 18s.

II.

THE IRRITABLE BLADDER: its Causes and Curative Treatment. Post 8vo. cloth, 4s. 6d.

DR. GIBB, M.R.C.P.

I.

ON DISEASES OF THE THROAT AND WINDPIPE, as reflected by the Laryngoscope. Second Edition. With 116 Engravings. Post 8vo. cloth, 10s. 6d.

II.

THE LARYNGOSCOPE IN DISEASES OF THE THROAT, with a Chapter on Rhinoscopy. Second Edition, enlarged, with Engravings. Crown 8vo., cloth, 5s.

MRS. GODFREY.

ON THE NATURE, PREVENTION, TREATMENT, AND CURE OF SPINAL CURVATURES and DEFORMITIES of the CHEST and LIMBS, without ARTIFICIAL SUPPORTS or any MECHANICAL APPLIANCES. Third Edition, Revised and Enlarged. 8vo. cloth 5s.

DR. GORDON M.D. C.B.

I.

ARMY HYGIENE. 8vo. cloth, 20s.

II.

CHINA, FROM A MEDICAL POINT OF VIEW, IN 1860 AND 1861; With a Chapter on Nagasaki as a Sanatorium. 8vo. cloth, 10s. 6d.

DR. GAIRDNER.

ON GOUT; its History, its Causes, and its Cure. Fourth Edition. Post 8vo. cloth, 8s. 6d.

DR. GRANVILLE, F.R.S.

THE MINERAL SPRINGS OF VICHY: their Efficacy in the Treatment of Gout, Indigestion, Gravel, &amp;c. 8vo. cloth, 3s.

ON SUDDEN DEATH. Post 8vo., 2s. 6d.

DR. GRAVES M.D., F.R.S.

STUDIES IN PHYSIOLOGY AND MEDICINE. Edited by Dr. Stokes. With Portrait and Memoir. 8vo. cloth, 14s.

DR. S. C. GRIFFITH, M.D.

ON DERMATOLOGY AND THE TREATMENT OF SKIN DISEASES BY MEANS OF HERBS, IN PLACE OF ARSENIC AND MERCURY. Fcap. 8vo. cloth, 3s.

MR. GRIFFITHS.

CHEMISTRY OF THE FOUR SEASONS—Spring, Summer, Autumn, Winter. Illustrated with Engravings on Wood. Second Edition. Foolscep 8vo. cloth, 7s. 6d.

DR. GULLY.

THE SIMPLE TREATMENT OF DISEASE; deduced from the Methods of Expectancy and Revulsion. 18mo. cloth, 4s.

DR. GUY AND DR. JOHN HARLEY.

HOOPER'S PHYSICIAN'S VADE-MECUM; OR, MANUAL OF THE PRINCIPLES AND PRACTICE OF PHYSIC. Seventh Edition, considerably enlarged, and rewritten. Foolscep 8vo. cloth, 12s. 6d.

GUY'S HOSPITAL REPORTS. Third Series. Vols. I. to XII., 8vo., 7s. 6d. each.

DR. HABERSHON, F.R.C.P.

ON DISEASES OF THE ABDOMEN, comprising those of the Stomach and other Parts of the Alimentary Canal, Oesophagus, Stomach, Cæcum, Intestines, and Peritoneum. Second Edition, with Plates. 8vo. cloth, 14s.

ON THE INJURIOUS EFFECTS OF MERCURY IN THE TREATMENT OF DISEASE. Post 8vo. cloth, 3s. 6d.

DR. C. RADCLYFFE HALL.

TORQUAY IN ITS MEDICAL ASPECT AS A RESORT FOR PULMONARY INVALIDS. Post 8vo. cloth, 5s.

DR. MARSHALL HALL, F.R.S.

PRONE AND POSTURAL RESPIRATION IN DROWNING AND OTHER FORMS OF APNŒA OR SUSPENDED RESPIRATION. Post 8vo. cloth, 5s.

PRACTICAL OBSERVATIONS AND SUGGESTIONS IN MEDICINE. Second Series. Post 8vo. cloth, 8s. 6d.

MR. HARDWICH.

A MANUAL OF PHOTOGRAPHIC CHEMISTRY. With Engravings. Seventh Edition. Foolscap 8vo. cloth, 7s. 6d.

DR. J. BOWER HARRISON, M.D., M.R.C.P.

I.  
LETTERS TO A YOUNG PRACTITIONER ON THE DISEASES OF CHILDREN. Foolscap 8vo. cloth, 3s.

II.  
ON THE CONTAMINATION OF WATER BY THE POISON OF LEAD, and its Effects on the Human Body. Foolscap 8vo. cloth, 3s. 6d.

DR. HARTWIG.

I.  
ON SEA BATHING AND SEA AIR. Second Edition. Fcap. 8vo., 2s. 6d.

II.  
ON THE PHYSICAL EDUCATION OF CHILDREN. Fcap. 8vo., 2s. 6d.

DR. A. H. HASSALL.

I.  
THE URINE, IN HEALTH AND DISEASE; being an Explanation of the Composition of the Urine, and of the Pathology and Treatment of Urinary and Renal Disorders. Second Edition. With 79 Engravings (23 Coloured). Post 8vo. cloth, 12s. 6d.

II.  
THE MICROSCOPIC ANATOMY OF THE HUMAN BODY, IN HEALTH AND DISEASE. Illustrated with Several Hundred Drawings in Colour. Two vols. 8vo. cloth, £1. 10s.

MR. ALFRED HAVILAND, M.R.C.S.

CLIMATE, WEATHER, AND DISEASE; being a Sketch of the Opinions of the most celebrated Ancient and Modern Writers with regard to the Influence of Climate and Weather in producing Disease. With Four coloured Engravings. 8vo. cloth, 7s.

DR. HEADLAND, M.D., F.R.C.P.

I.  
ON THE ACTION OF MEDICINES IN THE SYSTEM. Fourth Edition. 8vo. cloth, 14s.

II.  
A MEDICAL HANDBOOK; comprehending such Information on Medical and Sanitary Subjects as is desirable in Educated Persons. Second Thousand. Foolscap 8vo. cloth, 5s.

DR. HEALE.

I.  
A TREATISE ON THE PHYSIOLOGICAL ANATOMY OF THE LUNGS. With Engravings. 8vo. cloth, 8s.

II.  
. A TREATISE ON VITAL CAUSES. 8vo. cloth, 9s.



MR. CHRISTOPHER HEATH, F.R.C.S.

I.

**PRACTICAL ANATOMY:** a Manual of Dissections. With numerous Engravings. Fcap. 8vo. cloth, 10s. 6d.

II.

**A MANUAL OF MINOR SURGERY AND BANDAGING, FOR THE USE OF HOUSE-SURGEONS, DRESSERS, AND JUNIOR PRACTITIONERS.** With Illustrations. Third Edition. Fcap. 8vo. cloth, 5s.

MR. HIGGINBOTTOM, F.R.S., F.R.C.S.E.

**A PRACTICAL ESSAY ON THE USE OF THE NITRATE OF SILVER IN THE TREATMENT OF INFLAMMATION, WOUNDS, AND ULCERS.** Third Edition, 8vo. cloth, 6s.

DR. HINDS.

**THE HARMONIES OF PHYSICAL SCIENCE IN RELATION TO THE HIGHER SENTIMENTS;** with Observations on Medical Studies, and on the Moral and Scientific Relations of Medical Life. Post 8vo. cloth, 4s.

MR. J. A. HINGESTON, M.R.C.S.

**TOPICS OF THE DAY, MEDICAL, SOCIAL, AND SCIENTIFIC.** Crown 8vo. cloth, 7s. 6d.

DR. HODGES.

**THE NATURE, PATHOLOGY, AND TREATMENT OF PUPERAL CONVULSIONS.** Crown 8vo. cloth, 3s.

DR. DECIMUS HODGSON.

**THE PROSTATE GLAND, AND ITS ENLARGEMENT IN OLD AGE.** With 12 Plates. Royal 8vo. cloth, 6s.

MR. JABEZ HOGG.

**A MANUAL OF OPHTHALMOSCOPIC SURGERY;** being a Practical Treatise on the Use of the Ophthalmoscope in Diseases of the Eye. Third Edition. With Coloured Plates. 8vo. cloth, 10s. 6d.

MR. LUTHER HOLDEN, F.R.C.S.

I.

**HUMAN OSTEOLOGY:** with Plates, showing the Attachments of the Muscles. Third Edition. 8vo. cloth, 16s.

II.

**A MANUAL OF THE DISSECTION OF THE HUMAN BODY.** With Engravings on Wood. Second Edition. 8vo. cloth, 16s.



MR BARNARD HOLT, F.R.C.S.

ON THE IMMEDIATE TREATMENT OF STRICTURE OF  
THE URETHRA. Second Edition, Enlarged. 8vo. cloth, 3s.

DR. W. CHARLES HOOD.

SUGGESTIONS FOR THE FUTURE PROVISION OF CRIMI-  
NAL LUNATICS. 8vo. cloth, 5s. 6d.

• DR. P. HOOD.

THE SUCCESSFUL TREATMENT OF SCARLET FEVER;  
also, OBSERVATIONS ON THE PATHOLOGY AND TREATMENT OF  
CROWING INSPIRATIONS OF INFANTS. Post 8vo. cloth, 5s.

MR. JOHN HORSLEY.

A CATECHISM OF CHEMICAL PHILOSOPHY; being a Familiar  
Exposition of the Principles of Chemistry and Physics. With Engravings on Wood.  
Designed for the Use of Schools and Private Teachers. Post 8vo. cloth, 6s. 6d.

MR. LUKE HOWARD, F.R.S.

ESSAY ON THE MODIFICATIONS OF CLOUDS. Third Edition,  
by W. D. and E. HOWARD. With 6 Lithographic Plates, from Pictures by Kenyon.  
4to. cloth, 10s. 6d.

DR. HAMILTON HOWE, M.D.

A THEORETICAL INQUIRY INTO THE PHYSICAL CAUSE  
OF EPIDEMIC DISEASES. Accompanied with Tables. 8vo. cloth, 7s.

DR. HUFELAND.

THE ART OF PROLONGING LIFE. Second Edition. Edited  
by ERASMUS WILSON, F.R.S. Foolscep 8vo., 2s. 6d.

MR. W. CURTIS HUGMAN, F.R.C.S.

ON HIP-JOINT DISEASE; with reference especially to Treatment  
by Mechanical Means for the Relief of Contraction and Deformity of the Affected Limb.  
With Plates. Re-issue, enlarged. 8vo. cloth, 3s. 6d.

MR. HULKE, F.R.C.S.


A PRACTICAL TREATISE ON THE USE OF THE  
OPHTHALMOSCOPE. Being the Jacksonian Prize Essay for 1859. Royal 8vo.  
cloth, 8s.

DR. HENRY HUNT.

## ON HEARTBURN AND INDIGESTION. 8vo. cloth, 5s.

PROFESSOR HUXLEY, F.R.S.

LECTURES ON THE ELEMENTS OF COMPARATIVE  
ANATOMY.—ON CLASSIFICATION AND THE SKULL. With 111 Illus-  
trations. 8vo. cloth, 10s. 6d.



MR. JONATHAN HUTCHINSON, F.R.C.S.

A CLINICAL MEMOIR ON CERTAIN DISEASES OF THE EYE AND EAR, CONSEQUENT ON INHERITED SYPHILIS; with an appended Chapter of Commentaries on the Transmission of Syphilis from Parent to Offspring, and its more remote Consequences. With Plates and Woodcuts, 8vo. cloth, 9s.

DR. INMAN, M.R.C.P.

I.  
ON MYALGIA: ITS NATURE, CAUSES, AND TREATMENT; being a Treatise on Painful and other Affections of the Muscular System. Second Edition. 8vo. cloth, 9s.

II.  
FOUNDATION FOR A NEW THEORY AND PRACTICE OF MEDICINE. Second Edition. Crown 8vo. cloth, 10s.



DR. JAGO, M.D. OXON., A.B. CANTAB.

ENTOPTICS, WITH ITS USES IN PHYSIOLOGY AND MEDICINE. With 54 Engravings. Crown 8vo. cloth, 5s.

MR. J. H. JAMES, F.R.C.S.

PRACTICAL OBSERVATIONS ON THE OPERATIONS FOR STRANGULATED HERNIA. 8vo. cloth, 5s.

DR. PROSSER JAMES, M.D.

SORE-THROAT: ITS NATURE, VARIETIES, AND TREATMENT; including the Use of the LARYNGOSCOPE as an Aid to Diagnosis. Second Edition, with numerous Engravings. Post 8vo. cloth, 5s.

DR. HANDFIELD JONES, M.B., F.R.C.P.

CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS. Post 8vo. cloth, 10s. 6d.

DR. H. BENCE JONES, M.D., F.R.S.

LECTURES ON SOME OF THE APPLICATIONS OF CHEMISTRY AND MECHANICS TO PATHOLOGY AND THERAPEUTICS. 8vo. cloth, 12s.

DR. HANDFIELD JONES, F.R.S., & DR. EDWARD H. SIEVEKING.

A MANUAL OF PATHOLOGICAL ANATOMY. Illustrated with numerous Engravings on Wood. Foolscap 8vo. cloth, 12s. 6d.



DR. JAMES JONES, M.D., M.R.C.P.

ON THE USE OF PERCHLORIDE OF IRON AND OTHER CHALYBEATE SALTS IN THE TREATMENT OF CONSUMPTION. Crown 8vo. cloth, 3s. 6d.

MR. WHARTON JONES, F.R.S.

I.  
A MANUAL OF THE PRINCIPLES AND PRACTICE OF  
OPHTHALMIC MEDICINE AND SURGERY; with Nine Coloured Plates and  
173 Wood Engravings. Third Edition, thoroughly revised. Foolscap 8vo. cloth, 12s. 6d.

II.  
THE WISDOM AND BENEFICENCE OF THE ALMIGHTY,  
AS DISPLAYED IN THE SENSE OF VISION. Actonian Prize Essay. With  
Illustrations on Steel and Wood. Foolscap 8vo. cloth, 4s. 6d.

III.  
DEFECTS OF SIGHT AND HEARING: their Nature, Causes, Pre-  
vention, and General Management. Second Edition, with Engravings. Fcap. 8vo. 2s. 6d.

IV.  
A CATECHISM OF THE MEDICINE AND SURGERY OF  
THE EYE AND EAR. For the Clinical Use of Hospital Students. Fcap. 8vo. 2s. 6d.

V.  
A CATECHISM OF THE PHYSIOLOGY AND PHILOSOPHY  
OF BODY, SENSE, AND MIND. For Use in Schools and Colleges. Fcap. 8vo.,  
2s. 6d.

MR. FURNEAUX JORDAN, M.R.C.S.

AN INTRODUCTION TO CLINICAL SURGERY; WITH A  
Method of Investigating and Reporting Surgical Cases. Fcap. 8vo. cloth, 5s.

MR. JUDD.

A PRACTICAL TREATISE ON URETHRITIS AND SYPHI-  
LIS: including Observations on the Power of the Menstruous Fluid, and of the Dis-  
charge from Leucorrhœa and Sores to produce Urethritis: with a variety of Examples,  
Experiments, Remedies, and Cures. 8vo. cloth, £1. 5s.

DR. LAENNEC.

A MANUAL OF AUSCULTATION AND PERCUSSION. Trans-  
lated and Edited by J. B. SHARPE, M.R.C.S. 3s.

DR. LANE, M.A.

HYDROPATHY; OR, HYGIENIC MEDICINE. An Explanatory  
Essay. Second Edition. Post 8vo. cloth, 5s.

MR. LAWRENCE, F.R.S.

I.  
LECTURES ON SURGERY. 8vo. cloth, 16s.

II.  
A TREATISE ON RUPTURES. The Fifth Edition, considerably  
enlarged. 8vo. cloth, 16s.

DR. LEARED, M.R.C.P.

IMPERFECT DIGESTION: ITS CAUSES AND TREATMENT.  
Fourth Edition. Foolscap 8vo. cloth, 4s.



DR. EDWIN LEE.

I.  
THE EFFECT OF CLIMATE ON TUBERCULOUS DISEASE,  
with Notices of the chief Foreign Places of Winter Resort. Small 8vo. cloth, 4s. 6d.

II.  
THE WATERING PLACES OF ENGLAND, CONSIDERED  
with Reference to their Medical Topography. Fourth Edition. Fcap. 8vo. cloth, 7s. 6d.

III.  
THE PRINCIPAL BATHS OF FRANCE. Fourth Edition.  
Fcap. 8vo. cloth, 3s. 6d.

IV.  
THE BATHS OF GERMANY. Fourth Edition. Post 8vo. cloth, 7s.

V.  
THE BATHS OF SWITZERLAND. 12mo. cloth, 3s. 6d.

VI.  
HOMŒOPATHY AND HYDROPATHY IMPARTIALLY AP-  
PRECIATED. Fourth Edition. Post 8vo. cloth, 3s.


MR. HENRY LEE, F.R.C.S.

I.  
ON SYPHILIS. Second Edition. With Coloured Plates. 8vo. cloth, 10s.

II.  
ON DISEASES OF THE VEINS, HÆMORRHOIDAL TUMOURS,  
AND OTHER AFFECTIONS OF THE RECTUM. Second Edition. 8vo. cloth, 8s.

DR. ROBERT LEE, F.R.S.

I.  
CONSULTATIONS IN MIDWIFERY. Foolscap 8vo. cloth, 4s. 6d.

II.  
A TREATISE ON THE SPECULUM; with Three Hundred Cases.   
8vo. cloth, 4s. 6d.

III.  
CLINICAL REPORTS OF OVARIAN AND UTERINE DIS-  
EASES, with Commentaries. Foolscap 8vo. cloth, 6s. 6d.

IV.  
CLINICAL MIDWIFERY; comprising the Histories of 545 Cases of  
Difficult, Prematural, and Complicated Labour, with Commentaries. Second Edition.  
Foolscap 8vo. cloth, 5s.

DR. LEISHMAN, M.D., F.F.P.S.

THE MECHANISM OF PARTURITION: An Essay, Historical and  
Critical. With Engravings. 8vo. cloth, 5s.

MR. LISTON, F.R.S.

PRACTICAL SURGERY. Fourth Edition. 8vo. cloth, 22s.

MR. H. W. LOBB, L.S.A., M.R.C.S.E.

ON SOME OF THE MORE OBSCURE FORMS OF NERVOUS  
AFFECTIONS, THEIR PATHOLOGY AND TREATMENT. Re-issue,  
with the Chapter on Galvanism entirely Re-written. With Engravings. 8vo. cloth, 8s.

DR. LOGAN, M.D., M.R.C.P. LOND.

ON OBSTINATE DISEASES OF THE SKIN. Fcap. 8vo. cloth, 2s. 6d.

## LONDON HOSPITAL.

CLINICAL LECTURES AND REPORTS BY THE MEDICAL  
AND SURGICAL STAFF. With Illustrations. Vols. I. to III. 8vo. cloth, 7s. 6d.

## LONDON MEDICAL SOCIETY OF OBSERVATION.

WHAT TO OBSERVE AT THE BED-SIDE, AND AFTER  
DEATH. Published by Authority. Second Edition. Foolscep 8vo. cloth, 4s. 6d.

## MR. M'CLELLAND, F.L.S., F.G.S.

THE MEDICAL TOPOGRAPHY, OR CLIMATE AND SOILS,  
OF BENGAL AND THE N. W. PROVINCES. Post 8vo. cloth, 4s. 6d.

## DR. MACLACHLAN, M.D., F.R.C.P.L.

THE DISEASES AND INFIRMITIES OF ADVANCED LIFE.  
8vo. cloth, 16s.

## DR. A. C. MACLEOD, M.R.C.P. LOND.

ACHOLIC DISEASES; comprising Jaundice, Diarrhoea, Dysentery,  
and Cholera. Post 8vo. cloth, 5s. 6d.

## DR. GEORGE H. B. MACLEOD, F.R.C.S.E.

I.  
OUTLINES OF SURGICAL DIAGNOSIS. 8vo. cloth, 12s. 6d.

II.  
NOTES ON THE SURGERY OF THE CRIMEAN WAR; with  
REMARKS on GUN-SHOT WOUNDS. 8vo. cloth, 10s. 6d.

## MR. JOSEPH MACLISE, F.R.C.S.

I.  
SURGICAL ANATOMY. A Series of Dissections, illustrating the Prin-  
cipal Regions of the Human Body.  
The Second Edition, imperial folio, cloth, £3. 12s.; half-morocco, £4. 4s.

II.  
ON DISLOCATIONS AND FRACTURES. This Work is Uniform  
with the Author's "Surgical Anatomy;" each Fasciculus contains Four beautifully  
executed Lithographic Drawings. Imperial folio, cloth, £2. 10s.; half-morocco, £2. 17s.

## MR. MACNAMARA.

ON DISEASES OF THE EYE; referring principally to those Affections  
requiring the aid of the Ophthalmoscope for their Diagnosis. With coloured plates.  
8vo. cloth, 10s. 6d.

## DR. MONICOLL, M.R.C.P.

A HAND-BOOK FOR SOUTHPORT, MEDICAL & GENERAL;  
with Copious Notices of the Natural History of the District. Second Edition. Post 8vo.  
cloth, 3s. 6d.

## DR. MARCET, F.R.S.

I.  
ON THE COMPOSITION OF FOOD, AND HOW IT IS  
ADULTERATED; with Practical Directions for its Analysis. 8vo. cloth, 6s. 6d.

II.  
ON CHRONIC ALCOHOLIC INTOXICATION; with an INQUIRY  
INTO THE INFLUENCE OF THE ABUSE OF ALCOHOL AS A PRE-  
DISPOSING CAUSE OF DISEASE. Second Edition, much enlarged. Foolscep  
8vo. cloth, 4s. 6d.

DR. J. MACPHERSON, M.D.

**CHOLERA IN ITS HOME;** with a Sketch of the Pathology and Treatment of the Disease. Crown vo. cloth, 5s.

DR. MARKHAM.

I.  
**DISEASES OF THE HEART: THEIR PATHOLOGY, DIAGNOSIS, AND TREATMENT.** Second Edition. Post 8vo. cloth, 6s.

II.  
**SKODA ON AUSCULTATION AND PERCUSSION.** Post 8vo. cloth, 6s.

III.  
**BLEEDING AND CHANGE IN TYPE OF DISEASES.** Gulstonian Lectures for 1864. Crown 8vo. 2s. 6d.

SIR RANALD MARTIN, K.C.B., F.R.S.

**INFLUENCE OF TROPICAL CLIMATES IN PRODUCING THE ACUTE ENDEMIC DISEASES OF EUROPEANS;** including Practical Observations on their Chronic Sequelæ under the Influences of the Climate of Europe. Second Edition, much enlarged. 8vo. cloth, 20s.

DR. MASSY.

**ON THE EXAMINATION OF RECRUITS;** intended for the Use of Young Medical Officers on Entering the Army. 8vo. cloth, 5s.

MR. C. F. MAUNDER, F.R.C.S.

**OPERATIVE SURGERY.** With 158 Engravings. Post 8vo. 6s.

DR. MAYNE, M.D., LL.D.

I.  
**AN EXPOSITORY LEXICON OF THE TERMS, ANCIENT AND MODERN, IN MEDICAL AND GENERAL SCIENCE.** 8vo. cloth, £2. 10s.

II.  
**A MEDICAL VOCABULARY;** or, an Explanation of all Names, Synonymes, Terms, and Phrases used in Medicine and the relative branches of Medical Science. Second Edition. Fcap. 8vo. cloth, 8s. 6d.

DR. MERYON, M.D., F.R.C.P.

**PATHOLOGICAL AND PRACTICAL RESEARCHES ON THE VARIOUS FORMS OF PARALYSIS.** 8vo. cloth, 6s.

DR. MILLINGEN.

**ON THE TREATMENT AND MANAGEMENT OF THE INSANE;** with Considerations on Public and Private Lunatic Asylums. 18mo. cloth, 4s. 6d.

DR. W. J. MOORE, M.D.

I.  
**HEALTH IN THE TROPICS;** or, Sanitary Art applied to Europeans in India. 8vo. cloth, 9s.

II.  
**A MANUAL OF THE DISEASES OF INDIA.** Fcap. 8vo. cloth, 5s.

PROFESSOR MULDER, UTRECHT.

THE CHEMISTRY OF WINE. Edited by H. BENCE JONES, M.D.,  
F.R.S. Fcap. 8vo. cloth, 6s.

DR. W. MURRAY, M.D., M.R.C.P.

EMOTIONAL DISORDERS OF THE SYMPATHETIC SYSTEM OF NERVES. Crown 8vo. cloth, 3s. 6d.

DR. MUSHET, M.B., M.R.C.P.

ON APOPLEXY, AND ALLIED AFFECTIONS OF THE BRAIN. 8vo. cloth, 7s.

MR. NAYLER, F.R.C.S.

ON THE DISEASES OF THE SKIN. With Plates. 8vo. cloth, 10s. 6d.

DR. BIRKBECK NEVINS.

THE PRESCRIBER'S ANALYSIS OF THE BRITISH PHARMACOPEIA. Third Edition, enlarged to 295 pp. 32mo. cloth, 3s. 6d.

DR. THOS. NICHOLSON, M.D.

ON YELLOW FEVER; comprising the History of that Disease as it appeared in the Island of Antigua. Fcap. 8vo. cloth, 2s. 6d.

DR. NOAD, PH.D., F.R.S.

THE INDUCTION COIL, being a Popular Explanation of the Electrical Principles on which it is constructed. Second Edition. With Engravings. Fcap. 8vo. cloth, 3s.

DR. NOBLE.

THE HUMAN MIND IN ITS RELATIONS WITH THE BRAIN AND NERVOUS SYSTEM. Post 8vo. cloth, 4s. 6d.

MR. NUNNELEY, F.R.C.S.E.

ON THE ORGANS OF VISION: THEIR ANATOMY AND PHYSIOLOGY. With Plates, 8vo. cloth, 15s.

A TREATISE ON THE NATURE, CAUSES, AND TREATMENT OF ERYSIPELAS. 8vo. cloth, 10s. 6d.

MR. LANGSTON PARKER.

THE MODERN TREATMENT OF SYPHILITIC DISEASES, both Primary and Secondary; comprising the Treatment of Constitutional and Confirmed Syphilis, by a safe and successful Method. Fourth Edition, 8vo. cloth, 10s.



**MESSRS. CHURCHILL & SONS' PUBLICATIONS.**

**DR. PARKIN, M.D., F.R.C.S.**

**I.**  
**THE ANTIDOTAL TREATMENT AND PREVENTION OF**  
**THE EPIDEMIC CHOLERA.** Third Edition. 8vo. cloth, 7s. 6d.

**II.**  
**THE CAUSATION AND PREVENTION OF DISEASE; with**  
the Laws regulating the Extrication of Malaria from the Surface, and its Diffusion in the  
surrounding Air. 8vo. cloth, 5s.

**MR. JAMES PART, F.R.C.S.**

**THE MEDICAL AND SURGICAL POCKET CASE BOOK,**  
for the Registration of important Cases in Private Practice, and to assist the Student of  
Hospital Practice. Second Edition. 2s. 6d.

**DR. PAVY, M.D., F.R.S., F.R.C.P.**

**DIABETES: RESEARCHES ON ITS NATURE AND TREAT-**  
**MENT.** 8vo. cloth, 8s. 6d.

**DR. PEACOCK, M.D., F.R.C.P.**

**I.**  
**ON MALFORMATIONS OF THE HUMAN HEART.** With  
Original Cases and Illustrations. Second Edition. With 8 Plates. 8vo. cloth, 10s.

**II.**  
**ON SOME OF THE CAUSES AND EFFECTS OF VALVULAR**  
**DISEASE OF THE HEART.** With Engravings. 8vo. cloth, 5s.

**DR. W. H. PEARSE, M.D. EDIN.**

**NOTES ON HEALTH IN CALCUTTA AND BRITISH**  
**EMIGRANT SHIPS,** including Ventilation, Diet, and Disease. Fcap. 3vo. 2s.

**DR. PEET, M.D., F.R.C.P.**

**THE PRINCIPLES AND PRACTICE OF MEDICINE;**  
Designed chiefly for Students of Indian Medical Colleges. 8vo. cloth, 16s.

**DR. PEREIRA, F.R.S.**

**SELECTA E PRÆSCRIPTIS.** Fourteenth Edition. 24mo. cloth, 5s.

**DR. PICKFORD.**

**HYGIENE; or, Health as Depending upon the Conditions of the Atmo-**  
sphere, Food and Drinks, Motion and Rest, Sleep and Wakefulness, Secretions, Excre-  
tions, and Retentions, Mental Emotions, Clothing, Bathing, &c. Vol. I. 8vo. cloth, 9s.

**PROFESSOR PIRRIE, F.R.S.E.**

**THE PRINCIPLES AND PRACTICE OF SURGERY.** With  
numerous Engravings on Wood. Second Edition. 8vo. cloth, 24s.

**PROFESSOR PIRRIE & DR. KEITH.**

**ACUPRESSURE: an excellent Method of arresting Surgical Hæmorrhage**  
and of accelerating the healing of Wounds. With Engravings. 8vo. cloth, 5s.

**PROFESSORS PLATTNER & MUSPRATT.**

**THE USE OF THE BLOWPIPE IN THE EXAMINATION OF**  
**MINERALS, ORES, AND OTHER METALLIC COMBINATIONS.** Illustrated  
by numerous Engravings on Wood. Third Edition. 8vo. cloth, 10s. 6d.

DR. HENRY F. A. PRATT, M.D., M.R.C.P.

I.  
**THE GENEALOGY OF CREATION**, newly Translated from the Unpointed Hebrew Text of the Book of Genesis, showing the General Scientific Accuracy of the Cosmogony of Moses and the Philosophy of Creation. 8vo. cloth, 14s.

II.  
**ON ECCENTRIC AND CENTRIC FORCE: A New Theory of Projection.** With Engravings. 8vo. cloth, 10s.

III.  
**ON ORBITAL MOTION: The Outlines of a System of Physical Astronomy.** With Diagrams. 8vo. cloth, 7s. 6d.

IV.  
**ASTRONOMICAL INVESTIGATIONS.** The Cosmical Relations of the Revolution of the Lunar Apsides. Oceanic Tides. With Engravings. 8vo. cloth, 5s.

V.  
**THE ORACLES OF GOD: An Attempt at a Re-interpretation.** Part I. The Revealed Cosmos. 8vo. cloth, 10s.

~~~~~  
**THE PRESCRIBER'S PHARMACOPŒIA**; containing all the Medicines in the British Pharmacopœia, arranged in Classes according to their Action, with their Composition and Doses. By a Practising Physician. Fifth Edition. 32mo. cloth, 2s. 6d.; roan tuck (for the pocket), 3s. 6d.

~~~~~  
DR. JOHN ROWLISON PRETTY.

**AIDS DURING LABOUR**, including the Administration of Chloroform, the Management of Placenta and Post-partum Hæmorrhage. Fcap. 8vo. cloth, 4s. 6d.

MR. P. C. PRICE, F.R.C.S.

**AN ESSAY ON EXCISION OF THE KNEE-JOINT.** With Coloured Plates. With Memoir of the Author and Notes by Henry Smith, F.R.C.S. Royal 8vo. cloth, 14s.

~~~~~  
DR. PRIESTLEY.

**LECTURES ON THE DEVELOPMENT OF THE GRAVID UTERUS.** 8vo. cloth, 5s. 6d.

~~~~~  
DR. RADCLIFFE, F.R.C.P.L.

**LECTURES ON EPILEPTIC, SPASMODIC, NEURALGIC, AND PARALYTIC DISORDERS OF THE NERVOUS SYSTEM**, delivered at the Royal College of Physicians in London. Post 8vo. cloth, 7s. 6d.

~~~~~  
MR. RAINEY.

**ON THE MODE OF FORMATION OF SHELLS OF ANIMALS, OF BONE, AND OF SEVERAL OTHER STRUCTURES**, by a Process of Molecular Coalescence, Demonstrable in certain Artificially-formed Products. Fcap. 8vo. cloth, 4s. 6d.

~~~~~  
DR. F. H. RAMSBOTHAM.

**THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY.** Illustrated with One Hundred and Twenty Plates on Steel and Wood; forming one thick handsome volume. Fourth Edition. 8vo. cloth, 22s.



DR. RAMSBOTHAM.

PRACTICAL OBSERVATIONS ON MIDWIFERY, with a Selection  
of Cases. Second Edition. 8vo. cloth, 12s.

PROFESSOR REDWOOD, PH.D.

A SUPPLEMENT TO THE PHARMACOPŒIA: A concise but  
comprehensive Dispensatory, and Manual of Facts and Formulæ, for the use of Practi-  
tioners in Medicine and Pharmacy. Third Edition. 8vo. cloth, 22s.

DR. DU BOIS REYMOND.


ANIMAL ELECTRICITY; Edited by H. BENGE JONES, M.D., F.R.S.  
With Fifty Engravings on Wood. Foolscap 8vo. cloth, 6s.

DR. REYNOLDS, M.D. LOND.

I.  
EPILEPSY: ITS SYMPTOMS, TREATMENT, AND RELATION  
TO OTHER CHRONIC CONVULSIVE DISEASES. 8vo. cloth, 10s.

II.  
THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL  
CORD, AND THEIR APPENDAGES. 8vo. cloth, 8s.

DR. B. W. RICHARDSON.

I.  
ON THE CAUSE OF THE COAGULATION OF THE BLOOD.   
Being the ASTLEY COOPER PRIZE ESSAY for 1856. With a Practical Appendix.  
8vo. cloth, 16s.

II.  
† THE HYGIENIC TREATMENT OF PULMONARY CONSUMP-  
TION. 8vo. cloth, 5s. 6d.

DR. RITCHIE, M.D.

ON OVARIAN PHYSIOLOGY AND PATHOLOGY. With  
Engravings. 8vo. cloth, 6s.

DR. WILLIAM ROBERTS, M.D., F.R.C.P.

AN ESSAY ON WASTING PALSY; being a Systematic Treatise on  
the Disease hitherto described as ATROPHIE MUSCULAIRE PROGRESSIVE.  
With Four Plates. 8vo. cloth, 5s.

DR. ROUTH.

INFANT FEEDING, AND ITS INFLUENCE ON LIFE;  
Or, the Causes and Prevention of Infant Mortality. Second Edition. Fcap. 8vo. cloth, 6s.

DR. W. H. ROBERTSON.

I.  
THE NATURE AND TREATMENT OF GOUT. 8vo. cloth, 10s. 6d.

II.  
A TREATISE ON DIET AND REGIMEN. Fourth Edition. 2 vols  
12s. post 8vo. cloth.

DR. ROWE.

NERVOUS DISEASES, LIVER AND STOMACH COMPLAINTS, LOW SPIRITS, INDIGESTION, GOUT, ASTHMA, AND DISORDERS PRODUCED BY TROPICAL CLIMATES. With Cases. Sixteenth Edition. Fcap. 8vo. 2s. 6d.

DR. ROYLE, F.R.S., AND DR. HEADLAND, M.D.

A MANUAL OF MATERIA MEDICA AND THERAPEUTICS. With numerous Engravings on Wood. Fourth Edition. Fcap. 8vo. cloth, 12s. 6d.

DR. RYAN, M.D.

INFANTICIDE: ITS LAW, PREVALENCE, PREVENTION, AND HISTORY. 8vo. cloth, 5s.

ST. BARTHOLOMEW'S HOSPITAL

A DESCRIPTIVE CATALOGUE OF THE ANATOMICAL MUSEUM. Vol. I. (1846), Vol. II. (1851), Vol. III. (1862), 8vo. cloth, 5s. each.

ST. GEORGE'S HOSPITAL REPORTS. Vol. I. 8vo. 7s. 6d.

MR. T. P. SALT, BIRMINGHAM.

ON DEFORMITIES AND DEBILITIES OF THE LOWER EXTREMITIES, AND THE MECHANICAL TREATMENT EMPLOYED IN THE PROMOTION OF THEIR CURE. With numerous Plates. 8vo. cloth, 15s.

ON RUPTURE: ITS CAUSES, MANAGEMENT, AND CURE, and the various Mechanical Contrivances employed for its Relief. With Engravings. Post 8vo. cloth, 3s.

DR. SALTER, F.R.S.

ON ASTHMA: its Pathology, Causes, Consequences, and Treatment. 8vo. cloth, 10s.

DR. SANKEY, M.D. LOND.

LECTURES ON MENTAL DISEASES. 8vo. cloth, 8s.

DR. SANSOM M.D. LOND.

CHLOROFORM: ITS ACTION AND ADMINISTRATION. A Handbook. With Engravings. Crown 8vo. cloth, 5s.

THE ARREST AND PREVENTION OF CHOLERA; being a Guide to the Antiseptic Treatment. Fcap. 8vo. cloth, 2s. 6d.

MR. SAVORY.

A COMPENDIUM OF DOMESTIC MEDICINE, AND COMPANION TO THE MEDICINE CHEST; intended as a Source of Easy Reference for Clergymen, and for Families residing at a Distance from Professional Assistance. Seventh Edition. 12mo. cloth, 5s.

DR. SCHACHT.

THE MICROSCOPE, AND ITS APPLICATION TO VEGETABLE ANATOMY AND PHYSIOLOGY. Edited by FREDERICK CURREY, M.A. Fcap. 8vo. cloth, 6s.

DR. SCORESBY-JACKSON, M.D., F.R.S.E.

MEDICAL CLIMATOLOGY; or, a Topographical and Meteorological Description of the Localities resorted to in Winter and Summer by Invalids of various classes both at Home and Abroad. With an Isothermal Chart. Post 8vo. cloth, 12s.

DR. SEMPLE.

ON COUGH: its Causes, Varieties, and Treatment. With some practical Remarks on the Use of the Stethoscope as an aid to Diagnosis. Post 8vo. cloth, 4s. 6d.

DR. SEYMOUR.

I.  
ILLUSTRATIONS OF SOME OF THE PRINCIPAL DISEASES OF THE OVARIA: their Symptoms and Treatment; to which are prefixed Observations on the Structure and Functions of those parts in the Human Being and in Animals. On India paper. Folio, 16s.

II.  
THE NATURE AND TREATMENT OF DROPSY; considered especially in reference to the Diseases of the Internal Organs of the Body, which most commonly produce it. 8vo. 5s.

DR. SHAPTER, M.D., F.R.C.P.

THE CLIMATE OF THE SOUTH OF DEVON, AND ITS INFLUENCE UPON HEALTH. Second Edition, with Maps. 8vo. cloth, 10s. 6d.

MR. SHAW, M.R.C.S.

THE MEDICAL REMEMBRANCER; OR, BOOK OF EMERGENCIES. Fourth Edition. Edited, with Additions, by JONATHAN HUTCHINSON, F.R.C.S. 32mo. cloth, 2s. 6d.

DR. SHEA, M.D., B.A.

A MANUAL OF ANIMAL PHYSIOLOGY. With an Appendix of Questions for the B.A. London and other Examinations. With Engravings. Foolscap 8vo. cloth, 5s. 6d.

DR. SHRIMPTON.

CHOLERA: ITS SEAT, NATURE, AND TREATMENT. With Engravings. 8vo. cloth, 4s. 6d.

DR. SIBSON, F.R.S.

MEDICAL ANATOMY. With coloured Plates. Imperial folio. Fasciculi I. to VI. 5s. each.

DR. E. H. SIEVEKING.

ON EPILEPSY AND EPILEPTIFORM SEIZURES: their Causes, Pathology, and Treatment. Second Edition. Post 8vo. cloth, 10s. 6d.

DR. SIMMS.

A WINTER IN PARIS: being a few Experiences and Observations of French Medical and Sanitary Matters. Fcap. 8vo. cloth, 4s.

MR. SINCLAIR AND DR. JOHNSTON.

PRACTICAL MIDWIFERY: Comprising an Account of 13,748 Deliveries, which occurred in the Dublin Lying-in Hospital, during a period of Seven Years. 8vo. cloth, 10s.

DR. SIORDET, M.B.LOND., M.R.C.P.

MENTONE IN ITS MEDICAL ASPECT. Foolscap 8vo. cloth, 2s. 6d.

MR. ALFRED SMEE, F.R.S.

GENERAL DEBILITY AND DEFECTIVE NUTRITION; their Causes, Consequences, and Treatment. Second Edition. Fcap. 8vo. cloth, 3s. 6d.

DR. SMELLIE.

† **OBSTETRIC PLATES:** being a Selection from the more Important and Practical Illustrations contained in the Original Work. With Anatomical and Practical Directions. 8vo. cloth, 5s.

MR. HENRY SMITH, F.R.C.S.

I.  
**ON STRICTURE OF THE URETHRA.** 8vo. cloth, 7s. 6d.

II.  
**HÆMORRHOIDS AND PROLAPSUS OF THE RECTUM:**  
Their Pathology and Treatment, with especial reference to the use of Nitric Acid. Third Edition. Fcap. 8vo. cloth, 3s.

III.  
**THE SURGERY OF THE RECTUM.** Lettsomian Lectures. Fcap. 8vo. 2s. 6d.

DR. J. SMITH, M.D., F.R.C.S. EDIN.

**HANDBOOK OF DENTAL ANATOMY AND SURGERY, FOR**  
THE USE OF STUDENTS AND PRACTITIONERS. Fcap. 8vo. cloth, 3s. 6d.

DR. W. TYLER SMITH.

‡ **A MANUAL OF OBSTETRICS, THEORETICAL AND PRACTICAL.** Illustrated with 186 Engravings. Fcap. 8vo. cloth, 12s. 6d.

DR. SNOW.

**ON CHLOROFORM AND OTHER ANÆSTHETICS: THEIR**  
ACTION AND ADMINISTRATION. Edited, with a Memoir of the Author, by Benjamin W. Richardson, M.D. 8vo. cloth, 10s. 6d.

MR. J. VOSE SOLOMON, F.R.C.S.

**TENSION OF THE EYEBALL; GLAUCOMA:** some Account of the Operations practised in the 19th Century. 8vo. cloth, 4s.

DR. STANHOPE TEMPLEMAN SPEER.

**PATHOLOGICAL CHEMISTRY, IN ITS APPLICATION TO**  
THE PRACTICE OF MEDICINE. Translated from the French of MM. BECQUEREL and RODIER. 8vo. cloth, reduced to 8s.

MR. BALMANNO SQUIRE, M.B. LOND.

I.  
**COLOURED PHOTOGRAPHS OF SKIN DISEASES.** With Descriptive Letterpress. Series I. (12 Parts), 42s.; Series II. (6 Parts), 22s. 6d.; Series III. (6 Parts), 21s.

II.  
**A STUDENT'S ATLAS OF SKIN DISEASES.** (A Series of Chromolithographs, with Descriptive Letterpress). With cloth portfolio, 25s.

MR. PETER SQUIRE.

I.  
**A COMPANION TO THE BRITISH PHARMACOPŒIA.**  
Third Edition. 8vo. cloth, 8s. 6d.

II.  
**THE PHARMACOPŒIAS OF THIRTEEN OF THE LONDON**  
HOSPITALS, arranged in Groups for easy Reference and Comparison. 18mo. cloth,

## DR. STEGGALL.

## STUDENTS' BOOKS FOR EXAMINATION.

I.

A MEDICAL MANUAL FOR APOTHECARIES' HALL AND OTHER MEDICAL BOARDS. Twelfth Edition. 12mo. cloth, 10s.

II.

A MANUAL FOR THE COLLEGE OF SURGEONS; intended for the Use of Candidates for Examination and Practitioners. Second Edition. 12mo. cloth, 10s.

III.

GREGORY'S CONSPECTUS MEDICINÆ THEORETICÆ. The First Part, containing the Original Text, with an Ordo Verborum, and Literal Translation. 12mo. cloth, 10s.

IV.

THE FIRST FOUR BOOKS OF CELSUS; containing the Text, Ordo Verborum, and Translation. Second Edition. 12mo. cloth, 8s.

V.

FIRST LINES FOR CHEMISTS AND DRUGGISTS PREPARING FOR EXAMINATION AT THE PHARMACEUTICAL SOCIETY. Second Edition. 18mo. cloth, 3s. 6d.

## MR. STOWE, M.R.C.S.

A TOXICOLOGICAL CHART, exhibiting at one view the Symptoms, Treatment, and Mode of Detecting the various Poisons, Mineral, Vegetable, and Animal. To which are added, concise Directions for the Treatment of Suspended Animation. Twelfth Edition, revised. On Sheet, 2s.; mounted on Roller, 5s.

## MR. FRANCIS SUTTON, F.C.S.

A SYSTEMATIC HANDBOOK OF VOLUMETRIC ANALYSIS; or, the Quantitative Estimation of Chemical Substances by Measure. With Engravings. Post 8vo. cloth, 7s. 6d.

## DR. SWAYNE.

OBSTETRIC APHORISMS FOR THE USE OF STUDENTS COMMENCING MIDWIFERY PRACTICE. With Engravings on Wood. Third Edition. Fcap. 8vo. cloth, 3s. 6d.

## MR. TAMPLIN, F.R.C.S.E.

LATERAL CURVATURE OF THE SPINE: its Causes, Nature, and Treatment. 8vo. cloth, 4s.

## DR. ALEXANDER TAYLOR, F.R.S.E.

THE CLIMATE OF PAU; with a Description of the Watering Places of the Pyrenees, and of the Virtues of their respective Mineral Sources in Disease. Third Edition. Post 8vo. cloth, 7s.

## DR. ALFRED S. TAYLOR, F.R.S.

I.

THE PRINCIPLES AND PRACTICE OF MEDICAL JURISPRUDENCE. With 176 Wood Engravings. 8vo. cloth, 28s.

II.

A MANUAL OF MEDICAL JURISPRUDENCE. Eighth Edition. With Engravings. Fcap. 8vo. cloth, 12s. 6d.

III.

ON POISONS, in relation to MEDICAL JURISPRUDENCE AND MEDICINE. Second Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. TEALE.

ON AMPUTATION BY A LONG AND A SHORT RECTANGULAR FLAP. With Engravings on Wood. 8vo. cloth, 5s.

DR. THEOPHILUS THOMPSON, F.R.S.

CLINICAL LECTURES ON PULMONARY CONSUMPTION; with additional Chapters by E. SYMES THOMPSON, M.D. With Plates. 8vo. cloth, 7s. 6d.

DR. THOMAS.

THE MODERN PRACTICE OF PHYSIC; exhibiting the Symptoms, Causes, Morbid Appearances, and Treatment of the Diseases of all Climates. Eleventh Edition. Revised by ALGERNON FRAMPTON, M.D. 2 vols. 8vo. cloth, 28s.

MR. HENRY THOMPSON, F.R.C.S.

I.  
STRICTURE OF THE URETHRA; its Pathology and Treatment. The Jacksonian Prize Essay for 1852. With Plates. Second Edition. 8vo. cloth, 10s.

II.  
THE DISEASES OF THE PROSTATE; their Pathology and Treatment. Comprising a Dissertation "On the Healthy and Morbid Anatomy of the Prostate Gland;" being the Jacksonian Prize Essay for 1860. With Plates. Second Edition. 8vo. cloth, 10s.

III.  
PRACTICAL LITHOTOMY AND LITHOTRITY; or, An Inquiry into the best Modes of removing Stone from the Bladder. With numerous Engravings, 8vo. cloth, 9s.

DR. THUDICHUM.

I.  
A TREATISE ON THE PATHOLOGY OF THE URINE, Including a complete Guide to its Analysis. With Plates, 8vo. cloth, 14s.

II.  
A TREATISE ON GALL STONES: their Chemistry, Pathology, and Treatment. With Coloured Plates. 8vo. cloth, 10s.

DR. TILT.

I.  
ON UTERINE AND OVARIAN INFLAMMATION, AND ON THE PHYSIOLOGY AND DISEASES OF MENSTRUATION. Third Edition. 8vo. cloth, 12s.

II.  
A HANDBOOK OF UTERINE THERAPEUTICS, AND OF MODERN PATHOLOGY OF DISEASES OF WOMEN. Second Edition. Post 8vo. cloth, 6s.

III.  
THE CHANGE OF LIFE IN HEALTH AND DISEASE: a Practical Treatise on the Nervous and other Affections incidental to Women at the Decline of Life. Second Edition. 8vo. cloth, 6s.

DR. GODWIN TIMMS.

CONSUMPTION: its True Nature and Successful Treatment. Re-issue, enlarged. Crown 8vo. cloth, 10s.



DR. ROBERT B. TODD, F.R.S.

## CLINICAL LECTURES ON THE PRACTICE OF MEDICINE.

*New Edition, in one Volume, Edited by Dr. BEALE, 8vo. cloth, 18s.*ON CERTAIN DISEASES OF THE URINARY ORGANS, AND  
ON DROPSIES. Fcap. 8vo. cloth, 6s.

MR. TOMES, F.R.S.

A MANUAL OF DENTAL SURGERY. 'With 208 Engravings on  
Wood. Fcap. 8vo. cloth, 12s. 6d.

MR. JOSEPH TOYNBEE, F.R.S., F.R.C.S.

THE DISEASES OF THE EAR: THEIR NATURE, DIAG-  
NOSIS, AND TREATMENT. Illustrated with numerous Engravings on Wood.  
8vo. cloth, 15s.

DR. TURNBULL.

AN INQUIRY INTO THE CURABILITY OF CONSUMPTION,  
ITS PREVENTION, AND THE PROGRESS OF IMPROVEMENT IN THE  
TREATMENT. Third Edition. 8vo. cloth, 6s.A PRACTICAL TREATISE ON DISORDERS OF THE STOMACH  
with FERMENTATION; and on the Causes and Treatment of Indigestion, &c. 8vo.  
cloth, 6s.

DR. TWEEDIE, F.R.S.

CONTINUED FEVERS: THEIR DISTINCTIVE CHARACTERS,  
PATHOLOGY, AND TREATMENT. With Coloured Plates. 8vo. cloth, 12s.VESTIGES OF THE NATURAL HISTORY OF CREATION.  
Eleventh Edition. Illustrated with 106 Engravings on Wood. 8vo. cloth, 7s. 6d.

DR. UNDERWOOD.

TREATISE ON THE DISEASES OF CHILDREN. Tenth Edition,  
with Additions and Corrections by HENRY DAVIES, M.D. 8vo. cloth, 15s.

DR. UNGER.

BOTANICAL LETTERS. Translated by Dr. B. PAUL. Numerous  
Woodcuts. Post 8vo., 2s. 6d.

MR. WADE, F.R.C.S.

STRICTURE OF THE URETHRA, ITS COMPLICATIONS  
AND EFFECTS; a Practical Treatise on the Nature and Treatment of those  
Affections. Fourth Edition. 8vo. cloth, 7s. 6d.

DR. WALKER, M.B.LOND.

ON DIPHTHERIA AND DIPHTHERITIC DISEASES. Fcap.  
8vo. cloth, 3s.

DR. WALLER.

ELEMENTS OF PRACTICAL MIDWIFERY; or, Companion to  
the Lying-in Room. Fourth Edition, with Plates. Fcap. cloth, 4s. 6d.

MR. HAYNES WALTON, F.R.C.S.

SURGICAL DISEASES OF THE EYE. With Engravings on  
Wood. Second Edition. 8vo. cloth, 14s.

DR. WARING, M.D., M.R.C.P.LOND.

I.

A MANUAL OF PRACTICAL THERAPEUTICS. Second Edition,  
Revised and Enlarged. Fcap. 8vo. cloth, 12s. 6d.

II.

THE TROPICAL RESIDENT AT HOME. Letters addressed to  
Europeans returning from India and the Colonies on Subjects connected with their Health  
and General Welfare. Crown 8vo. cloth, 5s.

DR. WATERS, M.R.C.P.

I.

THE ANATOMY OF THE HUMAN LUNG. The Prize Essay  
to which the Fothergillian Gold Medal was awarded by the Medical Society of London.  
Post 8vo. cloth, 6s. 6d.

II.

RESEARCHES ON THE NATURE, PATHOLOGY, AND  
TREATMENT OF EMPHYSEMA OF THE LUNGS, AND ITS RELA-  
TIONS WITH OTHER DISEASES OF THE CHEST. With Engravings. 8vo.  
cloth, 5s.

DR. ALLAN WEBB, F.R.C.S.L.

THE SURGEON'S READY RULES FOR OPERATIONS IN  
SURGERY. Royal 8vo. cloth, 10s. 6d.

DR. WEBER.

A CLINICAL HAND-BOOK OF AUSCULTATION AND PER-  
CUSSION. Translated by JOHN COCKLE, M.D. 5s.

MR. SOELBERG WELLS, M.D., M.R.C.S.

ON LONG, SHORT, AND WEAK SIGHT, and their Treatment by  
the Scientific Use of Spectacles. Second Edition. With Plates. 8vo. cloth, 6s.

---

MR. T. SPENCER WELLS, F.R.C.S.

I.  
DISEASES OF THE OVARIES: THEIR DIAGNOSIS AND  
TREATMENT. Vol. I. 8vo. cloth, 9s.

II.  
SCALE OF MEDICINES WITH WHICH MERCHANT VES-  
SELS ARE TO BE FURNISHED, by command of the Privy Council for Trade;  
With Observations on the Means of Preserving the Health of Seamen, &c. &c.  
Seventh Thousand. Fcap. 8vo. cloth, 3s. 6d.

---

DR. WEST.

LECTURES ON THE DISEASES OF WOMEN. Third Edition.  
8vo. cloth, 16s.

---

DR. UVEDALE WEST.

ILLUSTRATIONS OF PUERPERAL DISEASES. Second Edi-  
tion, enlarged. Post 8vo. cloth, 5s.

---

MR. WHEELER.

HAND-BOOK OF ANATOMY FOR STUDENTS OF THE  
FINE ARTS. With Engravings on Wood. Fcap. 8vo., 2s. 6d.

---

DR. WHITEHEAD, F.R.C.S.

ON THE TRANSMISSION FROM PARENT TO OFFSPRING  
OF SOME FORMS OF DISEASE, AND OF MORBID TAINTS AND  
TENDENCIES. Second Edition. 8vo. cloth, 10s. 6d.

---

DR. WILLIAMS, F.R.S.

PRINCIPLES OF MEDICINE: An Elementary View of the Causes,  
Nature, Treatment, Diagnosis, and Prognosis, of Disease. With brief Remarks on  
Hygienics, or the Preservation of Health. The Third Edition. 8vo. cloth, 15s.

---

THE WIFE'S DOMAIN: the YOUNG COUPLE—the MOTHER—the NURSE  
—the NURSING. Post 8vo. cloth, 3s. 6d.

---

DR. J. HUME WILLIAMS.

UN SOUNDNESS OF MIND, IN ITS MEDICAL AND LEGAL  
CONSIDERATIONS. 8vo. cloth, 7s. 6d.

MR. ERASMUS WILSON, F.R.S.

I.  
**THE ANATOMIST'S VADE-MECUM: A SYSTEM OF HUMAN ANATOMY.** With numerous Illustrations on Wood. Eighth Edition. Foolscap 8vo. cloth, 12s. 6d.

II.  
**ON DISEASES OF THE SKIN: A SYSTEM OF CUTANEOUS MEDICINE.** Sixth Edition. 8vo. cloth, 18s.

**THE SAME WORK;** illustrated with finely executed Engravings on Steel, accurately coloured. 8vo. cloth, 36s.

III.  
**HEALTHY SKIN: A Treatise on the Management of the Skin and Hair** in relation to Health. Seventh Edition. Foolscap 8vo. 2s. 6d.

IV.  
**PORTRAITS OF DISEASES OF THE SKIN.** Folio. Fasciculi I. to XII., completing the Work. 20s. each. The Entire Work, half morocco, £13.

V.  
**THE STUDENT'S BOOK OF CUTANEOUS MEDICINE AND DISEASES OF THE SKIN.** Post 8vo. cloth, 8s. 6d.

VI.  
**ON SYPHILIS, CONSTITUTIONAL AND HEREDITARY; AND ON SYPHILITIC ERUPTIONS.** With Four Coloured Plates. 8vo. cloth, 16s.

VII.  
**A THREE WEEKS' SCAMPER THROUGH THE SPAS OF GERMANY AND BELGIUM,** with an Appendix on the Nature and Uses of Mineral Waters. Post 8vo. cloth, 6s. 6d.

VIII.  
**THE EASTERN OR TURKISH BATH:** its History, Revival in Britain, and Application to the Purposes of Health. Foolscap 8vo., 2s.

DR. G. C. WITTSTEIN.

**PRACTICAL PHARMACEUTICAL CHEMISTRY:** An Explanation of Chemical and Pharmaceutical Processes, with the Methods of Testing the Purity of the Preparations, deduced from Original Experiments. Translated from the Second German Edition, by STEPHEN DARBY. 18mo. cloth, 6s.

DR. HENRY J. WRIGHT.

**HEADACHES;** their Causes and their Cure. Fourth Edition. Fcap. 8vo. 2s. 6d.

DR. YEARSLEY, M.D., M.R.O.S.

I.  
**DEAFNESS PRACTICALLY ILLUSTRATED;** being an Exposition as to the Causes and Treatment of Diseases of the Ear. Sixth Edition. 8vo. cloth, 6s.

II.  
**ON THROAT AILMENTS, MORE ESPECIALLY IN THE ENLARGED TONSIL AND ELONGATED UVULA.** Eighth Edition. 8vo. cloth, 5s.



## **CHURCHILL'S SERIES OF MANUALS.**

Fcap. 8vo. cloth, 12s 6d. each.

"We here give Mr Churchill public thanks for the positive benefit conferred on the Medical Profession by the series of beautiful and cheap Manuals which bear his imprint."—*British and Foreign Medical Review.*

**AGGREGATE SALE, 141,000 COPIES.**

**ANATOMY.** With numerous Engravings. Eighth Edition. By ERASMUS WILSON, F.R.C.S., F.R.S.

**BOTANY.** With numerous Engravings. By ROBERT BENTLEY, F.L.S., Professor of Botany, King's College, and to the Pharmaceutical Society.

**CHEMISTRY.** With numerous Engravings. Ninth Edition. By GEORGE FOWNES, F.R.S., H. BENCE JONES, M.D., F.R.S., and A. W. HOFMANN, F.R.S.

**DENTAL SURGERY.** With numerous Engravings. By JOHN TONES, F.R.S.

**MATERIA MEDICA.** With numerous Engravings. Fourth Edition. By J. FORBES ROYLE, M.D., F.R.S., and FREDERICK W. HEADLAND, M.D., F.L.S.

**MEDICAL JURISPRUDENCE.** With numerous Engravings. Eighth Edition. By ALFRED SWAINE TAYLOR, M.D., F.R.S.

**PRACTICE OF MEDICINE.** Second Edition. By G. HILARO BARLOW, M.D., M.A.

**The MICROSCOPE and its REVELATIONS.** With numerous Plates and Engravings. Third Edition. By W. B. CARPENTER, M.D., F.R.S.

**NATURAL PHILOSOPHY.** With numerous Engravings. Fifth Edition. By GOLDING BIRD, M.D., M.A., F.R.S., and CHARLES BROOKE, M.B., M.A., F.R.S.

**OBSTETRICS.** With numerous Engravings. By W. TYLER SMITH, M.D., F.R.C.P.

**OPHTHALMIC MEDICINE and SURGERY.** With coloured Plates and Engravings on Wood. Third Edition. By T. WHARTON JONES, F.R.C.S., F.R.S.

**PATHOLOGICAL ANATOMY.** With numerous Engravings. By C. HANDFIELD JONES, M.B., F.R.S., and E. H. SIEVEKING, M.D., F.R.C.P.

**PHYSIOLOGY.** With numerous Engravings. Fourth Edition. By WILLIAM B. CARPENTER, M.D., F.R.S.

**POISONS.** Second Edition. By ALFRED SWAINE TAYLOR, M.D., F.R.S.

**PRACTICAL ANATOMY.** With numerous Engravings. (10s. 6d.) By CHRISTOPHER HEATH, F.R.C.S.

**PRACTICAL SURGERY.** With numerous Engravings. Fourth Edition. By Sir WILLIAM FERGUSSON, Bart., F.R.C.S., F.R.S.

**THERAPEUTICS.** Second Edition. By E. J. Waring, M.D., M.R.C.E.











